

THE THREE-COLOUR MODEL WITH DOMAIN WALL BOUNDARY CONDITIONS

HJALMAR ROSENGREN

*Dedicated to Dennis Stanton
on his 60th birthday*

ABSTRACT. We study the partition function for the three-colour model with domain wall boundary conditions. We express it in terms of certain special polynomials, which can be constructed recursively. Our method generalizes Kuperberg's proof of the alternating sign matrix theorem, replacing the six-vertex model used by Kuperberg with the eight-vertex-solid-on-solid model. As applications, we obtain some combinatorial results on three-colourings. We also conjecture an explicit formula for the free energy of the model.

1. INTRODUCTION

An *alternating sign matrix* is a square matrix with entries 1, -1 and 0, such that the non-zero entries in each row and column alternate in sign and add up to 1. Mills, Robbins and Rumsey [MRR] conjectured that the number of $n \times n$ alternating sign matrices is

$$A_n = \frac{1!4!7! \cdots (3n-2)!}{n!(n+1)! \cdots (2n-1)!}. \quad (1.1)$$

This conjecture was proved by Zeilberger [Z1]. Soon afterwards, a much simpler proof was found by Kuperberg [K], using the *six-vertex model* of statistical mechanics.

The six-vertex model is an example of an *ice model*, whose states can be identified with what we call ice graphs; see (2.2) below. Alternating sign matrices can be identified with ice graphs satisfying *domain wall boundary conditions*. For the six-vertex model, there is a closed formula for the corresponding partition function, the *Izergin–Korepin determinant* [I, ICK]. Kuperberg observed that in a special case, when all parameters of the model are cubic roots of unity, the partition function simply counts the number of states. He could then prove (1.1) by computing the corresponding limit of the determinant.

There is a natural two-parameter extension of the six-vertex model known as the eight-vertex-solid-on-solid (8VSOS) model. This model was introduced by Baxter [B2] as a tool for solving the eight-vertex model. It is elliptic, that is,

2000 *Mathematics Subject Classification.* 05A15, 33E05, 82B23.

Key words and phrases. Three-colour model, eight-vertex-solid-on-solid model, domain wall boundary conditions, partition function, alternating sign matrix.

Research supported by the Swedish Science Research Council (Vetenskapsrådet).

the Boltzmann weights are elliptic functions of the parameters. We stress that, in contrast to the eight-vertex model, the 8VSOS model is an ice model. In particular, for domain wall boundary conditions, its states can be identified with alternating sign matrices.

It is natural to ask what happens to the 8VSOS model under Kuperberg's specialization of the parameters. The answer turns out to be very satisfactory: it degenerates to the *three-colour model*. It is an observation of Lenard that ice graphs are in bijection with three-colourings of a square lattice, such that adjacent squares have distinct colour [L]. The three-colour model is defined by assigning independent weights to the three colours; the partition function is simply the corresponding generating function [B1]. Thus, one may hope that extending Kuperberg's work to the 8VSOS model would lead to new applications of statistical mechanics to combinatorics. That is precisely the object of the present study.

Apparently, the first step in this program is to generalize the Izergin–Korepin formula to the 8VSOS model. In a recent paper [R] we found such a generalization. In the trigonometric limit (which is intermediate between the six-vertex model and the general elliptic 8VSOS model), we used it to obtain a closed formula for a special case of the three-colour partition function, see (2.5). In the present paper, we consider the general case. Although there seems to be no very simple formula for the general three-colour partition function, we can express it in terms of certain special polynomials, which have remarkable properties and deserve further study. To obtain these results has not been straight-forward; in particular, we have not been able to work directly with the explicit formulas from [R]. Rather, we combine a simple consequence of those formulas with several further ideas.

The plan of the paper is as follows. In §2, we describe the three-colour model with domain wall boundary conditions, and its relation to ice graphs and alternating sign matrices. We refer to the states of the model as *three-coloured chessboards*. In §3, we state our main results in elementary form. §4 contains preliminaries on theta functions and the 8VSOS model. In §5, we prove our first main result, Theorem 3.1, which expresses the three-colour partition function Z_n^{3C} in terms of special polynomials q_n and r_n . This is a rather easy consequence of results in [R].

We then turn towards an alternative way of expressing Z_n^{3C} . In §6, we introduce a function Φ_n , which provides a one-parameter extension of Z_n^{3C} . For the six-vertex model, a similar function appears in Zeilberger's proof of the *refined* alternating sign matrix conjecture [Z2]. In §7, Φ_n is generalized to a multivariable theta function Ψ_n . These functions play a similar role as Schur polynomials do in Stroganov's proof of the alternating sign matrix theorem [St] (see also [O]). However, while the Schur polynomials are instances of the six-vertex partition function (with the crossing parameter a cubic root of unity), the function Ψ_n is *not* directly related to the 8VSOS partition function. The function Ψ_n has two important properties, the first being a determinant formula reminiscent of the Izergin–Korepin determinant. The second property is a symmetry with respect to inversion of each variable, meaning that it naturally lives on the Riemann sphere

rather than the torus. As a consequence, Ψ_n can be identified with a symmetric polynomial S_n . In §8, we specialize the variables in S_n , obtaining certain two-variable polynomials P_n and one-variable polynomials p_n . Using minor relations for the determinant defining Ψ_n , we obtain recursions for these polynomials, which can be used to derive many further properties. In §9, we return to the three-colour model, expressing Z_n^{3C} in terms of the polynomials p_n . This result has combinatorial consequences. For instance, for three-coloured chessboards of fixed size, we can compute the maximal and minimal possible number of squares of each colour, see Corollary 3.3.

Finally, in §10 we study the thermodynamic limit $n \rightarrow \infty$. Using non-rigorous arguments, we are led to an explicit formula for the free energy of the three-colour model with domain wall boundary conditions (Conjecture 3.14). From the viewpoint of physics, this is the main result of the paper. To prove Conjecture 3.14 rigorously is an interesting problem, which we expect to be rather difficult. For the six-vertex model with domain wall boundary conditions, a rigorous analysis has been done only recently [BF, BL1, BL2, BL3].

Note added in proof: Immediately after seeing an earlier version of the present paper, Vladimir Bazhanov and Vladimir Mangazeev sent me an interesting conjectured recursion for the polynomials p_n . They have also obtained a similar conjecture for P_n . These recursions give a much faster way of computing the three-colour partition function than those obtained in the present paper. Both conjectures can be found in [BM4], where the authors stress the resemblance to polynomials occurring in their analysis of the eight-vertex model [BM1, BM2, BM3]. It would be interesting to investigate if that relation can be made precise; for instance, that may give a link between the present work and the Painlevé VI equation.

Acknowledgements: This work was partly carried out while participating in the programme Discrete Integrable Systems at the Isaac Newton Institute for Mathematical Sciences, and I thank the institute and the programme organizers for their support. I would also like to thank Vladimir Bazhanov, Vladimir Mangazeev, Jacques Perk and, in particular, Don Zagier for their interest and for valuable comments and suggestions. Finally, I thank the anonymous referees for useful comments.

2. THREE-COLOURED CHESSBOARDS


We will refer to a state of the three-colour model with domain wall boundary conditions as a *three-coloured chessboard*. Fixing n , consider a chessboard of size $(n+1) \times (n+1)$. The squares will be labelled with three colours, which we identify with the three residue classes $0, 1, 2 \pmod 3$. We impose the following two rules. First, vertically or horizontally adjacent squares have distinct colours. Second, the north-west and south-east squares are labelled 0 and, as one proceeds away from these squares along the boundary, the colours increase with respect to the

cyclic order

$$0 < 1 < 2 < 0. \quad (2.1)$$

In particular, the north-east and south-west square are labelled $n \bmod 3$. As an example, when $n = 3$ there are seven three-coloured chessboards; these are displayed in Figure 1.

FIGURE 1. The seven three-coloured chessboards of size $n = 3$, where 0 is pictured as black, 1 as red and 2 as yellow.




We will briefly explain the bijections to alternating sign matrices and ice graphs mentioned in the introduction. Let $\begin{pmatrix} a & b \\ c & d \end{pmatrix}$ be a 2×2 -block of adjacent squares from a three-coloured chessboard, and choose representatives of the residue classes so that adjacent labels differ by exactly 1. For instance, if the original block is $\begin{pmatrix} 0 & 2 \\ 2 & 1 \end{pmatrix}$, we may choose the representatives $\begin{pmatrix} 3 & 2 \\ 2 & 1 \end{pmatrix}$. Having made such a choice, we contract each block to the number $(b + c - a - d)/2$, obtaining an $n \times n$ -matrix with entries $-1, 0, 1$. For instance, from the last chessboard in (??) we obtain

$$\begin{pmatrix} 0 & 1 & 0 \\ 1 & -1 & 1 \\ 0 & 1 & 0 \end{pmatrix}.$$

This gives a bijection from three-coloured chessboards to alternating sign matrices.

To obtain the bijection to ice graphs, we draw an arrow between any two adjacent squares in such a way that the larger label, with respect to the order (2.1), is to the right. For instance, the last chessboard in (??) corresponds to the arrow configuration



$$(2.2)$$

The result is a directed graph, where each internal vertex has two incoming and two outgoing edges. Considering vertices as oxygen atoms and incoming edges as hydrogen bonds, this can be viewed as a model for a two-dimensional sheet of ice.

We are interested in the generating function

$$\begin{aligned} Z_n^{3C}(t_0, t_1, t_2) &= \sum_{\text{chessboards}} \prod_{\text{squares}} t_{\text{colour}} \\ &= \sum_{k_0+k_1+k_2=(n+1)^2} N(k_0, k_1, k_2) t_0^{k_0} t_1^{k_1} t_2^{k_2}, \end{aligned} \quad (2.3)$$

where $N(k_0, k_1, k_2)$ denotes the number of three-coloured chessboards with exactly k_i squares of colour i . In physics terminology, Z_n^{3C} is the *partition function* of the three-colour model with domain wall boundary conditions.

It is difficult to study Z_n^{3C} by direct methods. Indeed, to compute $Z_n^{3C}(1, 1, 1) = A_n$ was an unsolved problem for more than a decade. In [R], we generalized that enumeration using the trigonometric 8VSOS model. Namely, we found a closed expression for $Z_n^{3C}(t_0, t_1, t_2)$ when

$$\frac{(t_0 t_1 + t_0 t_2 + t_1 t_2)^3}{(t_0 t_1 t_2)^2} = 27. \quad (2.4)$$

This surface can be parametrized by $t_i(\lambda, \mu) = \mu/(1 - \lambda\omega^i)^3$, where, as throughout the paper,

$$\omega = e^{2\pi i/3}.$$

By homogeneity, we may take $\mu = 1$. Then [R, Cor. 8.4],

$$\begin{aligned} Z_n^{3C} \left(\frac{1}{(1-\lambda)^3}, \frac{1}{(1-\lambda\omega)^3}, \frac{1}{(1-\lambda\omega^2)^3} \right) \\ = \frac{(1-\lambda\omega^2)^2(1-\lambda\omega^{n+1})^2}{(1-\lambda^3)^{n^2+2n+3}} (A_n(1+\omega^n\lambda^2) + (-1)^n C_n \omega^{2n}\lambda), \end{aligned} \quad (2.5)$$

where A_n is as in (1.1) and

$$C_n = \frac{2 \cdot 5 \cdots (3n-1)}{1 \cdot 4 \cdots (3n-2)} A_n \quad (2.6)$$

is the number of cyclically symmetric plane partitions in a cube of size n [A]. Although (2.4) is a strong restriction, it is sufficient for computing the moments

$$\sum_{k_0+k_1+k_2=(n+1)^2} N(k_0, k_1, k_2) k_i, \quad i = 0, 1, 2,$$

see [R, Cor. 8.5].

3. STATEMENT OF RESULTS

3.1. Polynomials q_n and r_n . In this Section, we state our main results. We begin with the following fact. As we will see in §5, it is a rather straight-forward consequence of results in [R].

Theorem 3.1. *Let*

$$T = T(t_0, t_1, t_2) = \frac{(t_0 t_1 + t_0 t_2 + t_1 t_2)^3}{(t_0 t_1 t_2)^2}. \quad (3.1)$$

Then, there exist polynomials q_n and r_n such that, for $n \equiv 0 \pmod{3}$, $Z_n^{3C}(t_0, t_1, t_2)$ equals

$$(-1)^{n+1} (t_0 t_1 t_2)^{\frac{n(n+2)}{3}} \left(\frac{(t_0 t_1 + t_0 t_2 + t_1 t_2)^2}{t_0 t_1 t_2} q_n(T) - 2^{\chi(n \text{ odd})} t_0 r_n(T) \right),$$

while for $n \equiv 1 \pmod{3}$ it equals

$$(-1)^{n+1} (t_0 t_1 t_2)^{\frac{n(n+2)}{3}} \left(\frac{t_0 t_1}{t_2} q_n(T) - 2^{\chi(n \text{ odd})} \frac{t_0 t_1 + t_0 t_2 + t_1 t_2}{t_2} r_n(T) \right)$$

and for $n \equiv 2 \pmod{3}$ it equals

$$(-1)^{n+1} (t_0 t_1 t_2)^{\frac{(n+1)^2}{3}} \left(q_n(T) - 2^{\chi(n \text{ odd})} \frac{t_0 t_1 + t_0 t_2 + t_1 t_2}{t_0 t_2} r_n(T) \right).$$

Here, $\chi(\text{true}) = 1$, $\chi(\text{false}) = 0$.

The first few instances of the polynomials q_n and r_n are given in Table 1.

Note that, in each case, Z_n^{3C} is symmetric in the two variables $t_{-n \pm 1 \pmod{3}}$. This is explained by the fact that reflection in the vertical (say) axis, followed by interchanging the colours $-n \pm 1 \pmod{3}$, defines an involution on three-coloured chessboards. Theorem 3.1 shows that Z_n^{3C} is very nearly symmetric in all three variables, being a linear combination of two symmetric polynomials, where the coefficients are polynomials in $t_{-n \pmod{3}}$ of low order. Moreover, the symmetric polynomials depend only on the second and third elementary symmetric polynomial, being independent of $t_0 + t_1 + t_2$.

TABLE 1. The polynomials q_n and r_n .

n	$q_n(x)$	$r_n(x)$
0	0	1
1	1	0
2	1	1
3	1	1
4	$x + 3$	$x - 3$
5	$x^2 - 4x + 6$	$x + 6$
6	$x^2 - 2x + 40$	$x^3 - 8x^2 + 20$
7	$x^4 - 10x^3 + 15x^2 + 100x + 50$	$x^3 + 75x - 50$

The following result seems much deeper than Theorem 3.1. We need considerable preparation for its proof, which is given in §9.2.

Theorem 3.2. *The polynomials q_n and r_n are monic. Moreover, their degrees are given by*

$$\begin{aligned} \deg(q_n) + 1 = \deg(r_n) &= \frac{n^2}{12}, & n &\equiv 0 \pmod{6}, \\ \deg(q_n) = \deg(r_n) + 1 &= \frac{n^2 - 1}{12}, & n &\equiv \pm 1 \pmod{6}, \\ \deg(q_n) = \deg(r_n) &= \frac{n^2 - 4}{12}, & n &\equiv \pm 2 \pmod{6}, \\ \deg(q_n) = \deg(r_n) &= \frac{n^2 - 9}{12}, & n &\equiv 3 \pmod{6}. \end{aligned}$$

As an application, we can determine the maximal and minimal number of squares of each colour. These bounds restrict the counting function N introduced in (2.3) to an equilateral triangle. We can also explicitly evaluate the restriction of N to the boundary. To formulate the result, we introduce some notation. Fixing n , let

$$\bar{N}(x, y, z) = \begin{cases} N(x, y, z), & n \equiv 0 \pmod{3}, \\ N(y, z, x), & n \equiv 1 \pmod{3}, \\ N(z, x, y), & n \equiv 2 \pmod{3}, \end{cases}$$

so that $\bar{N}(x, y, z) = \bar{N}(x, z, y)$. Moreover, let

$$M = \left\lfloor \frac{5n^2 + 8n + 11}{12} \right\rfloor = \begin{cases} (5n^2 + 8n)/12, & n \equiv 0, 2 \pmod{6}, \\ (5n^2 + 8n + 11)/12, & n \equiv 1 \pmod{6}, \\ (5n^2 + 8n + 3)/12, & n \equiv 3, 5 \pmod{6}, \\ (5n^2 + 8n + 8)/12, & n \equiv 4 \pmod{6}, \end{cases}$$

$$m = \begin{cases} (n^2 + 4n)/6, & n \equiv 0, 2 \pmod{6}, \\ (n^2 + 4n + 7)/6, & n \equiv 1 \pmod{6}, \\ (n^2 + 4n + 3)/6, & n \equiv 3, 5 \pmod{6}, \\ (n^2 + 4n + 4)/6, & n \equiv 4 \pmod{6}, \end{cases}$$

$$\varepsilon = \begin{cases} 1, & n \equiv 0, 2 \pmod{6}, \\ -2, & n \equiv 1 \pmod{6}, \\ 0, & n \equiv 3, 5 \pmod{6}, \\ -1, & n \equiv 4 \pmod{6}, \end{cases}$$

$$\delta = \left\lfloor \frac{n^2}{4} \right\rfloor = \begin{cases} (n^2 - 1)/4, & n \text{ odd}, \\ n^2/4, & n \text{ even}. \end{cases} \quad (3.2)$$

Note that $2M + m + \varepsilon = (n + 1)^2$. Let

$$\Delta = \{(x, y, z) \in \mathbb{Z}^3; x + y + z = (n + 1)^2, x \leq M + \varepsilon, y, z \leq M\}.$$

Then, Δ is an equilateral triangle, with δ lattice points on each side. We denote its corners by

$$P = (m + \varepsilon, M, M), \quad Q = (M + \varepsilon, m, M), \quad R = (M + \varepsilon, M, m).$$

Corollary 3.3. *The convex hull of the support of $\bar{N}$ is equal to Δ when n is odd and $\Delta \setminus \{P\}$ when n is even. In particular, the maximal number of squares of each of the colours $-n \pm 1 \pmod{3}$ is equal to M , and the minimal number of such squares is m . The maximal number of squares of colour $-n \pmod{3}$ is $M + \varepsilon$, and the minimal number is $m + \varepsilon$ if n is odd and $m + \varepsilon + 1$ if n is even. Moreover, the restriction of $\bar{N}$ to $\partial\Delta$ is given by*

$$\bar{N}\left(\frac{kP + (\delta - k)Q}{\delta}\right) = \bar{N}\left(\frac{kP + (\delta - k)R}{\delta}\right) = \begin{cases} \binom{\delta - 1}{k}, & n \text{ even,} \\ \binom{\delta}{k}, & n \text{ odd,} \end{cases}$$

$$\bar{N}\left(\frac{kQ + (\delta - k)R}{\delta}\right) = \begin{cases} \binom{\delta}{k}, & n \text{ even,} \\ \binom{\delta - 2}{k} + \binom{\delta - 2}{k - 2}, & n \text{ odd,} \end{cases}$$

where $0 \leq k \leq \delta$.

It is straight-forward to derive Corollary 3.3 from Theorem 3.2; some details are given in §9.2.

As we explain at the end of §5, it follows easily from Theorem 3.1 that the polynomials q_n , r_{2n} and $2r_{2n+1}$ have integer coefficients. However, the following fact is not obvious, see §9.3.

Proposition 3.4. *The polynomial r_{2n+1} has integer coefficients.*

This result has a simple combinatorial meaning. If $n \equiv 3$ or $5 \pmod{6}$, Theorem 3.1 expresses Z_n^{3C} as a sum of a symmetric polynomial and a polynomial with even coefficients. Thus, the function

$$(k_0, k_1, k_2) \mapsto N(k_0, k_1, k_2) \pmod{2}$$

is symmetric. Similarly, when $n \equiv 1 \pmod{6}$,

$$(k_0, k_1, k_2) \mapsto N(k_0, k_1, k_2 - 2) \pmod{2}$$

is symmetric. In the notation of Corollary 3.3, these facts can be stated as follows.

Corollary 3.5. *When n is odd, $\bar{N} \pmod{2}$ is invariant under the action of S_3 as the symmetry group of Δ .*

To illustrate Corollaries 3.3 and 3.5, we give two examples. When $n = 4$, the non-zero values of N are

$$(N(6+i, 6+j, 13-i-j))_{i,j=0}^4 = \begin{pmatrix} & & & & 1 \\ & & & 4 & 3 \\ & & 6 & 6 & 3 \\ & 4 & 6 & & 1 \\ 1 & 3 & 3 & 1 & \end{pmatrix}.$$

Since n is even, the lower right corner is missing from the support of N , and all boundary entries are binomial coefficients. When $n = 5$, the non-zero values are

$$(N(8+i, 20-i-j, 8+j))_{i,j=0}^6 = \begin{pmatrix} & & & & & & 1 \\ & & & & & 4 & 6 \\ & & & 7 & 18 & 15 \\ & & 8 & 12 & 36 & 20 \\ & 7 & 12 & 36 & 40 & 15 \\ 4 & 18 & 36 & 40 & 24 & 6 \\ 1 & 6 & 15 & 20 & 15 & 6 & 1 \end{pmatrix}.$$

In this case, the diagonal entries are numbers of the form $\binom{4}{k} + \binom{4}{k-2}$. Note the symmetric distribution of the odd entries, which is peculiar to the case of odd n .

3.2. Polynomials p_n and P_n . The polynomials q_n and r_n are closely related to a third class of polynomials, which we denote p_n . In the following result, which is proved in §9.2, we use the notation

$$\tilde{p}(x) = x^{\deg p} p(1/x); \quad (3.3)$$

that is, $\tilde{p}$ denotes the polynomial obtained from p by reversing the coefficients.

Theorem 3.6. *There exist polynomials p_n of degree $n(n+1)/2$ such that, for n odd,*

$$p_{n-1}(\zeta) - \zeta^{\frac{n+1}{2}} \tilde{p}_{n-1}(\zeta) = (1-\zeta)(\zeta^2 + 4\zeta + 1)^{\frac{n^2-1}{4}} \tilde{q}_n \left(\frac{\zeta(\zeta+1)^4}{2(\zeta^2 + 4\zeta + 1)^3} \right),$$

$$p_{n-1}(\zeta) - \zeta^{\frac{n-1}{2}} \tilde{p}_{n-1}(\zeta) = (1-\zeta)(1+\zeta)^3(\zeta^2 + 4\zeta + 1)^{\frac{n^2-9}{4}} \tilde{r}_n \left(\frac{\zeta(\zeta+1)^4}{2(\zeta^2 + 4\zeta + 1)^3} \right),$$

whereas for n even,

$$p_{n-1}(\zeta) - \zeta^{\frac{n+2}{2}} \tilde{p}_{n-1}(\zeta) = (1-\zeta)(\zeta^2 + 4\zeta + 1)^{\frac{n^2}{4}} \tilde{r}_n \left(\frac{\zeta(\zeta+1)^4}{2(\zeta^2 + 4\zeta + 1)^3} \right),$$

$$p_{n-1}(\zeta) - \zeta^{\frac{n}{2}} \tilde{p}_{n-1}(\zeta) = (1-\zeta^2)(\zeta^2 + 4\zeta + 1)^{\frac{n^2-4}{4}} \tilde{q}_n \left(\frac{\zeta(\zeta+1)^4}{2(\zeta^2 + 4\zeta + 1)^3} \right).$$

TABLE 2. The polynomials p_n .

n	$p_n(\zeta)$
-1	1
0	1
1	$3\zeta + 1$
2	$5\zeta^3 + 15\zeta^2 + 7\zeta + 1$
3	$\frac{1}{2}(35\zeta^6 + 231\zeta^5 + 504\zeta^4 + 398\zeta^3 + 147\zeta^2 + 27\zeta + 2)$
4	$\frac{1}{2}(63\zeta^{10} + 798\zeta^9 + 4122\zeta^8 + 11052\zeta^7 + 16310\zeta^6 + 13464\zeta^5 + 6636\zeta^4 + 2036\zeta^3 + 387\zeta^2 + 42\zeta + 2)$
5	$\frac{1}{2}(231\zeta^{15} + 4554\zeta^{14} + 39468\zeta^{13} + 196922\zeta^{12} + 622677\zeta^{11} + 1298446\zeta^{10} + 1816006\zeta^9 + 1726302\zeta^8 + 1140593\zeta^7 + 535478\zeta^6 + 181104\zeta^5 + 44134\zeta^4 + 7603\zeta^3 + 882\zeta^2 + 62\zeta + 2)$
6	$\frac{1}{8}(1716\zeta^{21} + 50193\zeta^{20} + 673530\zeta^{19} + 5484050\zeta^{18} + 30199260\zeta^{17} + 118703208\zeta^{16} + 342834244\zeta^{15} + 738954900\zeta^{14} + 1198556100\zeta^{13} + 1470762970\zeta^{12} + 1373623128\zeta^{11} + 984509064\zeta^{10} + 546520100\zeta^9 + 236837400\zeta^8 + 80476380\zeta^7 + 21422188\zeta^6 + 4430904\zeta^5 + 699405\zeta^4 + 81550\zeta^3 + 6630\zeta^2 + 336\zeta + 8)$

The shift in n is introduced for convenience. See Table 2 for the first few instances of the polynomials p_n .

To indicate the meaning of Theorem 3.6, we solve for p_n , obtaining for n even

$$p_n(\zeta) = \left((\zeta^2 + 4\zeta + 1)^{\frac{n(n+2)}{4}} \tilde{q}_{n+1} \left(\frac{\zeta(\zeta+1)^4}{2(\zeta^2 + 4\zeta + 1)^3} \right) - \zeta(1+\zeta)^3(\zeta^2 + 4\zeta + 1)^{\frac{(n-2)(n+4)}{4}} \tilde{r}_{n+1} \left(\frac{\zeta(\zeta+1)^4}{2(\zeta^2 + 4\zeta + 1)^3} \right) \right) \quad (3.4)$$

and for n odd

$$p_n(\zeta) = \left((\zeta^2 + 4\zeta + 1)^{\frac{(n+1)^2}{4}} \tilde{r}_{n+1} \left(\frac{\zeta(\zeta+1)^4}{2(\zeta^2 + 4\zeta + 1)^3} \right) - \zeta(1+\zeta)(\zeta^2 + 4\zeta + 1)^{\frac{(n-1)(n+3)}{4}} \tilde{q}_{n+1} \left(\frac{\zeta(\zeta+1)^4}{2(\zeta^2 + 4\zeta + 1)^3} \right) \right).$$

A priori, the right-hand sides are polynomials of degree $n(n+2)/2$, $(n+1)^2/2$, respectively. The degree bound $n(n+1)/2$ imposes relations between high coefficients of the polynomials q_n and r_n . In the notation of Corollary 3.3, this implies relations between values of $\bar{N}$ close to the boundary $\partial\Delta$; the details will not be worked out here.

An important property of the polynomials p_n is that they appear as solutions to certain linear equations.

Theorem 3.7. *Consider the polynomial equation*

$$AX - BY = C, \quad (3.5)$$

where

$$\begin{aligned} A &= (\zeta + 1)^2 p_{n-1}(\zeta), & B &= \zeta^{n+1} \tilde{p}_n(\zeta), \\ C &= (1 + 2\zeta)^{1+\chi(n \text{ even})} \left(1 + \frac{\zeta}{2}\right)^{1+\chi(n \text{ odd})} p_n(\zeta)^2. \end{aligned}$$

Then, this equation has a solution (X, Y) such that $X = p_{n+1}$ and $\deg Y = \deg A$.

Theorem 3.7 follows from Proposition 8.3 and Corollary 8.10. Although we have not been able to prove it, we believe that the polynomials A and B are always relatively prime. In fact, we believe that p_n is irreducible over $\mathbb{Q}$ for $n \geq 2$. Assuming that this is the case, (3.5) has a unique solution (X_0, Y_0) with $\deg Y_0 < \deg A$. The solutions with $\deg Y \leq \deg A$ then have the form $(X, Y) = (X_0 + \lambda B, Y_0 + \lambda A)$, $\lambda \in \mathbb{C}$. We can specify the solution $X = p_{n+1}$ in this space by declaring that the leading term is

$$2^{-\lfloor \frac{n+2}{2} \rfloor} \binom{2n+2}{n+1} \zeta^{n(n+1)/2},$$

see Proposition 8.9. Then, Theorem 3.7 gives a recursive procedure for constructing the polynomials p_n . This gives a comparatively fast method for computing the partition function Z_n^{3C} . (As was mentioned at the end of the Introduction, a much faster method has been suggested by Bazhanov and Mangazeev [BM4].)

The polynomials p_n have coefficients in $\mathbb{Z}/2^{\mathbb{Z}}$; see Corollary 9.8 for a more precise statement. Moreover, the following facts seem to be true.

Conjecture 3.8. *The polynomials p_n have positive coefficients.*

Conjecture 3.9. *The polynomials p_n are unimodal in the sense that, if*

$$p_n(\zeta) = \sum_{k=0}^{n(n+1)/2} a_k \zeta^k,$$

then, for some N ,


$$a_0 < a_1 < \cdots < a_{N-1} < a_N > a_{N+1} > \cdots > a_{n(n+1)/2}.$$

Using Theorem 3.7, we have verified Conjectures 3.8 and 3.9 up to $n = 16$. For these values, the maximal coefficient a_N occurs at

$$N = \begin{cases} n(n+2)/4, & n \text{ even}, \quad n \leq 16 \\ (n+1)^2/4, & n \text{ odd}, \quad n \leq 7 \\ (n-1)(n+3)/4, & n \text{ odd}, \quad 9 \leq n \leq 15. \end{cases}$$

Numerical experiments suggest that the zeroes of p_n form quite remarkable patterns, see Figure 2.

We have some partial results on the real zeroes, see §8.4 for proofs.

FIGURE 2. The 105 complex zeroes of p_{14} .

Conjecture 3.10. *The polynomial p_n has exactly $[(n+1)/2]$ real zeroes.*

Conjecture 3.11. *The polynomial p_n does not vanish in the interval $-2 < \zeta < -1/2$.*

Proposition 3.12. *Assume Conjecture 3.10. Then, all real zeroes of $p_n(\zeta)$ are simple, and contained in the interval $-1/2 < \zeta < 0$ if n is odd and in $\zeta < -2$ if n is even. Moreover, the real zeroes of p_{2n} interlace the real zeroes of $\tilde{p}_{2n+1}$, which in turn alternate left of the real zeroes of p_{2n+2} . In particular, Conjecture 3.10 implies Conjecture 3.11.*

The polynomials p_n can be embedded in a more general family of polynomials $P_n(x, \zeta)$, which are of degree n in x and degree $[n(n+2)/2]$ in ζ . To be precise,

$$P_n(1+2\zeta, \zeta) = (1+2\zeta)^{[\frac{n}{2}]} p_n(\zeta).$$

We give the first few instances of P_n in Table 3.

The polynomials P_n satisfy a three-term recursion of the form

$$A_n P_{n+1} = B_n P_n + C_n P_{n-1},$$

where $A_n = A_n(x, \zeta)$ and $C_n = C_n(x, \zeta)$ are quadratic polynomials in x , while $B_n = B_n(x, \zeta)$ is cubic in x , see Proposition 8.6. This is reminiscent of the recursion satisfied by orthogonal polynomials (where A_n and C_n are constants and B_n is linear). It also seems that P_n resemble orthogonal polynomials with respect to their zeroes. Indeed, the following fact is proved in §8.4.

Proposition 3.13. *Assume Conjecture 3.11 (or, in view of Proposition 3.12, Conjecture 3.10). Then, if $-2 < \zeta < -1/2$ and $\zeta \neq -1$, all zeroes of $P_n(x, \zeta)$ are simple and positive. Moreover, the zeroes of P_{n+1} interlace those of P_n .*

TABLE 3. The polynomials P_n .

n	$P_n(x, \zeta)$
0	1
1	$x + \zeta$
2	$\frac{1}{2} \left((\zeta + 2)(3\zeta + 1)x^2 + \zeta(\zeta + 3)(3\zeta + 1)x + \zeta^2(\zeta + 3)(2\zeta + 1) \right)$
3	$\frac{1}{2} \left((\zeta + 2)(5\zeta^3 + 15\zeta^2 + 7\zeta + 1)x^3 + \zeta(\zeta + 4)(5\zeta^3 + 15\zeta^2 + 7\zeta + 1)x^2 \right.$ $\left. + \zeta^2(4\zeta + 1)(\zeta^3 + 7\zeta^2 + 15\zeta + 5)x + \zeta^3(2\zeta + 1)(\zeta^3 + 7\zeta^2 + 15\zeta + 5) \right)$
4	$\frac{1}{8} \left((\zeta + 2)^2(35\zeta^6 + 231\zeta^5 + 504\zeta^4 + 398\zeta^3 + 147\zeta^2 + 27\zeta + 2)x^4 \right.$ $+ \zeta(\zeta + 5)(\zeta + 2)(35\zeta^6 + 231\zeta^5 + 504\zeta^4 + 398\zeta^3 + 147\zeta^2 + 27\zeta + 2)x^3$ $+ 3\zeta^2(10\zeta^8 + 139\zeta^7 + 790\zeta^6 + 2245\zeta^5$ $+ 3232\zeta^4 + 2245\zeta^3 + 790\zeta^2 + 139\zeta + 10)x^2$ $+ \zeta^3(2\zeta + 1)(5\zeta + 1)(2\zeta^6 + 27\zeta^5 + 147\zeta^4 + 398\zeta^3 + 504\zeta^2 + 231\zeta + 35)x$ $\left. + \zeta^4(2\zeta + 1)^2(2\zeta^6 + 27\zeta^5 + 147\zeta^4 + 398\zeta^3 + 504\zeta^2 + 231\zeta + 35) \right)$

Computer calculations suggest that, if $-2 < \zeta < -1$, the zeroes are in fact contained in the interval $x > 1$, while if $-1 < \zeta < -1/2$, they are contained in $0 < x < 1$.

3.3. Symmetric polynomials S_n . The key to the proof of most our results is that the polynomials P_n and p_n can be obtained as specializations of certain symmetric polynomials S_n of $2n + 1$ variables. To define them, we introduce the elementary polynomials

$$f(x) = (\zeta + 2)x - \zeta, \quad g(x) = (x - 1)(x - \zeta), \quad h(x) = x^2(x - (2\zeta + 1)), \quad (3.6)$$

and then let

$$F(x, y, z) = \frac{1}{(y - x)(z - x)(z - y)} \det \begin{pmatrix} f(x) & f(y) & f(z) \\ g(x) & g(y) & g(z) \\ h(x) & h(y) & h(z) \end{pmatrix} \quad (3.7)$$

$$= (\zeta + 2)xyz - \zeta(xy + yz + xz + x + y + z) + \zeta(2\zeta + 1),$$

$$G(x, y) = \frac{1}{y - x} \det \begin{pmatrix} f(x) & f(y) \\ h(x) & h(y) \end{pmatrix}$$

$$= (\zeta + 2)xy(x + y) - \zeta(x^2 + y^2) - 2(\zeta^2 + 3\zeta + 1)xy + \zeta(2\zeta + 1)(x + y). \quad (3.8)$$

In this notation,

$$S_n(x_1, \dots, x_n, y_1, \dots, y_n, z)$$

$$= \frac{\prod_{i,j=1}^n G(x_i, y_j)}{\prod_{1 \leq i < j \leq n} (x_j - x_i)(y_j - y_i)} \det_{1 \leq i, j \leq n} \left(\frac{F(x_i, y_j, z)}{G(x_i, y_j)} \right), \quad (3.9)$$

the dependence on ζ being suppressed from the notation. Though it is not apparent from this definition, S_n is a symmetric polynomial of all $2n + 1$ variables, see §7.2. Moreover,

$$\begin{aligned} S_n\left(x, \underbrace{2\zeta + 1, \dots, 2\zeta + 1}_n, \underbrace{\frac{\zeta}{\zeta + 2}, \dots, \frac{\zeta}{\zeta + 2}}_n\right) \\ = (-1)^{\binom{n+1}{2}} \zeta^{n^2} (1 + \zeta)^{n^2} (1 + 2\zeta)^{\left[\frac{(n-1)^2}{4}\right]} \left(1 + \frac{\zeta}{2}\right)^{\left[\frac{(n-1)^2}{4}\right] - n^2} P_n(x, \zeta), \end{aligned}$$

see Proposition 8.1.

We mention that the special case $\zeta = -2$ of S_n is an orthogonal character:

$$S_n(x_1, \dots, x_{2n+1}) \Big|_{\zeta=-2} = 2^{n^2} \chi_{(n,n,n-1,n-1,\dots,1,1,0)}^{\mathfrak{so}(4n+3)}(t_1, \dots, t_{2n+1}),$$

where $x_i = -(1 + t_i + t_i^{-1})$; see Theorem 7.17 for a more general result. For general ζ , S_n can probably be interpreted as an affine Lie algebra character; see Remark 7.4.

3.4. Thermodynamic limit. All results mentioned so far concern properties of the partition function for fixed n . However, from the viewpoint of statistical mechanics, the main problem is to investigate the asymptotics as $n \rightarrow \infty$. A particularly important quantity is the free energy per volume, which we identify with the limit

$$f(t_0, t_1, t_2) = \lim_{n \rightarrow \infty} \frac{\log Z_n^{3C}(t_0, t_1, t_2)}{n^2}. \quad (3.10)$$

We propose the following explicit expression for f . In §10 we present a formal derivation of this result; to give a rigorous proof is presumably quite difficult.

Conjecture 3.14. *Assume that the parameters t_i are all positive, and let ζ be any positive solution of*

$$T = \frac{(t_0 t_1 + t_0 t_2 + t_1 t_2)^3}{(t_0 t_1 t_2)^2} = \frac{2(\zeta^2 + 4\zeta + 1)^3}{\zeta(\zeta + 1)^4}. \quad (3.11)$$

Then, the free energy per volume is given by

$$f(t_0, t_1, t_2) = \frac{1}{3} \log(t_0 t_1 t_2) + \log \left(\frac{(\zeta + 2)^{\frac{3}{4}} (2\zeta + 1)^{\frac{3}{4}}}{2^{\frac{2}{3}} \zeta^{\frac{1}{12}} (\zeta + 1)^{\frac{4}{3}}} \right). \quad (3.12)$$

Note that, by the arithmetic-geometric inequality, $T \geq 27$. Using that

$$\frac{2(\zeta^2 + 4\zeta + 1)^3}{\zeta(\zeta + 1)^4} - 27 = \frac{(\zeta - 1)^4 (\zeta + 2)(2\zeta + 1)}{\zeta(\zeta + 1)^4}, \quad (3.13)$$

it is easy to check that (3.11) always has two positive solutions, except in the case $T = 27$, when $\zeta = 1$. Moreover, the two solutions are related by $\zeta \mapsto \zeta^{-1}$, which does not change the right-hand side of (3.12).

As an example, when $t_0 = t_1 = t_2 = \zeta = 1$, (3.12) gives

$$f(1, 1, 1) = \log \left(\frac{3\sqrt{3}}{4} \right),$$

which agrees with the known asymptotics [BF]

$$Z_n^{3C}(1, 1, 1) = A_n \sim C n^{-\frac{5}{36}} \left(\frac{3\sqrt{3}}{4} \right)^{n^2}, \quad n \rightarrow \infty. \quad (3.14)$$

One should compare Conjecture 3.14 with Baxter's result for *periodic* boundary conditions [B1]. In that case, one still has

$$f(t_0, t_1, t_2) = \frac{1}{3} \log(t_0 t_1 t_2) + \log W,$$

with W a function only of T . Baxter gives the formula

$$W^2 = \frac{64(1 - 9t^2)^{\frac{2}{3}}}{27(1 + t)^3(1 - 3t)},$$

where t is the unique solution of

$$\frac{T}{27} = \frac{(1 - 3t^2)^3}{1 - 9t^2}$$

such that $0 \leq t < 1/3$. It is straight-forward to check that $t = (\zeta - 1)/3(\zeta + 1)$, where ζ is the unique solution of (3.11) such that $\zeta \geq 1$. This gives

$$W = W_{\text{per}} = \frac{2^{\frac{5}{3}} \zeta^{\frac{1}{3}} (\zeta + 1)^{\frac{4}{3}}}{(2\zeta + 1)^{\frac{3}{2}}},$$

which is manifestly different from our conjectured formula

$$W_{\text{DWBC}} = \frac{(\zeta + 2)^{\frac{3}{4}} (2\zeta + 1)^{\frac{3}{4}}}{2^{\frac{2}{3}} \zeta^{\frac{1}{12}} (\zeta + 1)^{\frac{4}{3}}}$$

for domain wall boundary conditions. Note the intriguing relation

$$W_{\text{DWBC}}(\zeta) = \frac{2}{\sqrt{W_{\text{per}}(\zeta) W_{\text{per}}(1/\zeta)}}.$$

4. PRELIMINARIES

4.1. Theta functions. We will work on the multiplicative torus $\mathbb{C}^*/\{z = pz\}$, where $\mathbb{C}^* = \mathbb{C} \setminus \{0\}$ and $0 < |p| < 1$. We introduce the theta function

$$\theta(x; p) = \prod_{j=0}^{\infty} (1 - p^j x)(1 - p^{j+1}/x).$$

We often employ condensed notation such as

$$\theta(a_1, \dots, a_n; p) = \theta(a_1; p) \cdots \theta(a_n; p),$$

$$\theta(xy^\pm; p) = \theta(xy; p)\theta(xy^{-1}; p).$$

The most fundamental relations for the theta function are

$$\theta(px; p) = \theta(x^{-1}; p) = -x^{-1}\theta(x; p),$$

together with the addition formula

$$\theta(xz^\pm, yw^\pm; p) - \theta(xw^\pm, yz^\pm; p) = \frac{y}{z} \theta(xy^\pm, zw^\pm; p). \quad (4.1)$$

Other elementary identities that we will use include

$$\theta(x^2; p^2) = \theta(\pm x; p) = \theta(\pm x, \pm px; p^2),$$

from which one can derive

$$\theta(-1, \pm p; p^2) = 2, \quad (4.2a)$$

$$\theta(-\omega, \pm p\omega; p^2) = -\omega^2, \quad (4.2b)$$

where $\omega = e^{2\pi i/3}$.

We will use the following terminology from [RSc].

Definition 4.1. *A holomorphic function on $\mathbb{C}^*$ is called an A_{n-1} theta function of nome p and norm t if it satisfies*

$$f(px) = \frac{(-1)^n}{tx^n} f(x).$$

It is called a BC_n theta function of nome p if

$$f(px) = \frac{1}{p^n x^{2n+1}} f(x), \quad f(x^{-1}) = -x^{-1} f(x).$$

Finally, it is called a D_n theta function of nome p if

$$f(px) = \frac{1}{p^{n-1} x^{2n-2}}, \quad f(x^{-1}) = f(x).$$

Lemma 4.2 ([RSc, Lemma 3.2]). *A function f is an A_{n-1} theta function of nome p and norm t if and only if it can be factored as*

$$f(x) = C\theta(a_1x, \dots, a_nx; p),$$

where $C \in \mathbb{C}$ and $a_i \in \mathbb{C}^$ with $a_1 \cdots a_n = t$. It is a BC_n theta function of nome p if and only if*

$$f(x) = C\theta(x, \pm\sqrt{p}x; p)\theta(a_1x^\pm, \dots, a_{n-1}x^\pm; p),$$

where $C \in \mathbb{C}$ and $a_i \in \mathbb{C}^$. Finally, it is a D_n theta function of nome p if and only if*

$$f(x) = C\theta(a_1x^\pm, \dots, a_{n-1}x^\pm; p),$$

where $C \in \mathbb{C}$ and $a_i \in \mathbb{C}^$.*

The terminology is motivated by Macdonald's theory of affine root systems [M1]. In each case, the Macdonald identity for the affine root system R is equivalent to evaluating a determinant $\det(f_i(x_j))$, where f_i runs through a basis in the space of R theta functions, see [RSc, Proposition 6.1]. We will need two special cases, which are both classical theta function identities. For $R = A_1$, the Macdonald identity can be written

$$\theta(ax, bx; p) = \frac{1}{\theta(p; p^2)} (\theta(-pa/b, -abx^2; p^2) - bx\theta(-a/b, -pabx^2; p^2)); \quad (4.3)$$

this can also be obtained from (4.1). The Macdonald identity for BC_1 is Watson's quintuple product [W]

$$\theta(x, \pm\sqrt{p}x; p) = \frac{(p^3; p^3)_\infty}{(p; p)_\infty} (\theta(-px^3; p^3) - x\theta(-px^{-3}; p^3)), \quad (4.4)$$

where

$$(p; p)_\infty = \prod_{j=1}^{\infty} (1 - p^j). \quad (4.5)$$

Finally, we recall some classical facts on uniformization. The Riemann surface $S = \mathbb{C}^*/\{z = pz = z^{-1}\}$ is a sphere, the analytic automorphisms $\xi : S \rightarrow \mathbb{C} \cup \{\infty\}$ being given by

$$\xi(z) = C \frac{\theta(az^\pm; p)}{\theta(bz^\pm; p)},$$

where $C \in \mathbb{C}^*$ and a, b are distinct in $\mathbb{C}^*/\{z = pz = z^{-1}\}$. Accordingly, any D_{n+1} theta function f can be written

$$f(z) = \theta(bz^\pm; p)^n P(\xi(z)),$$

where P is a polynomial of degree at most n . We will refer to the passage from f to P as *uniformization*. When f is factored as in Lemma 4.2, the explicit uniformization is

$$\begin{aligned} \theta(a_1 z^\pm, \dots, a_n z^\pm; p) &= \frac{a_1 \cdots a_n \theta(ba_1^\pm, \dots, ba_n^\pm; p)}{C^n a^n \theta(ba^\pm; p)^n} \\ &\quad \times \theta(bz^\pm; p)^n (\xi(z) - \xi(a_1)) \cdots (\xi(z) - \xi(a_n)). \end{aligned} \quad (4.6)$$

This follows immediately from the case $n = 1$, which is equivalent to (4.1).

4.2. The 8VSOS model. The 8VSOS model is an ice model, so for domain wall boundary conditions states can be identified with three-coloured chessboards. Fixing the size of the chessboards to $(n+1) \times (n+1)$, the model depends on parameters

$$x_1, \dots, x_n, y_1, \dots, y_n, \lambda, p \in \mathbb{C}^*, \quad |p| < 1.$$

In general, there is also a crossing parameter q , but for our purposes it can be fixed to a cubic root of unity.

We will assign a weight to each 2×2 -block of adjacent squares. These blocks can be viewed as entries of an $n \times n$ matrix, and are given coordinates $1 \leq i, j \leq n$ in a standard way. A block $\begin{pmatrix} a & b \\ c & d \end{pmatrix}$ with coordinates (i, j) is then given the weight $R_{d-c, c-a}^{b-a, d-b}(\lambda \omega^a, x_i/y_j)$, where

$$\begin{aligned} R_{++}^{++}(\lambda, u) &= R_{--}^{--}(\lambda, u) = \frac{\theta(\omega u; p)}{\theta(u; p)}, \\ R_{+-}^{+-}(\lambda, u) &= \frac{\theta(u, \omega \lambda; p)}{\theta(\omega, \lambda; p)}, & R_{-+}^{-+}(\lambda, u) &= \omega \frac{\theta(u, \omega^2 \lambda; p)}{\theta(\omega, \lambda; p)}, \\ R_{-+}^{-+}(\lambda, u) &= \frac{\theta(\lambda u; p)}{\theta(\lambda; p)}, & R_{+-}^{+-}(\lambda, u) &= u \frac{\theta(\lambda/u; p)}{\theta(\lambda; p)}. \end{aligned}$$

Here, $\pm$ is a short-hand for $\pm 1 \bmod 3$.

The partition function is now defined as

$$Z_n^{8\text{VSOS}}(x_1, \dots, x_n; y_1, \dots, y_n; \lambda, p) = \sum_{\text{chessboards}} \prod_{\text{blocks}} \text{weight}(\text{block}).$$

This differs slightly from the normalization used in [R]. We have

$$Z_n^{8\text{VSOS}}(x; y; \lambda, p) = (y_1 \dots y_n)^{-n} \tilde{Z}_n(x; y; \lambda),$$

where $\tilde{Z}_n$ is as in [R, §7].

In the specialization $x_i \equiv \omega$, $y_i \equiv 1$, the 8VSOS model reduces to the three-colour model with parameters

$$t_i = \frac{1}{\theta(\lambda \omega^i; p)^3}. \quad (4.7)$$

More precisely, in the case of domain wall boundary conditions,

$$\begin{aligned} Z_n^{3C} \left(\frac{1}{\theta(\lambda; p)^3}, \frac{1}{\theta(\lambda \omega; p)^3}, \frac{1}{\theta(\lambda \omega^2; p)^3} \right) \\ = \omega^{n(n+1)} \frac{\theta(\lambda \omega^2, \lambda \omega^{n+1}; p)^2}{\theta(\lambda \omega^n; p) \theta(\lambda^3; p^3)^{n^2+2n+2}} Z_n^{8\text{VSOS}}(\omega, \dots, \omega; 1, \dots, 1; \lambda, p). \end{aligned} \quad (4.8)$$

See [R, §8] for the trigonometric case $p = 0$; the discussion there carries over *verbatim* to general p .

We need to mention the recursion

$$\begin{aligned} Z_n^{8\text{VSOS}}(x_1, \dots, x_n; y_1, \dots, y_n; \lambda, p) \Big|_{\omega x_1 = y_1} \\ = \omega^{n+1} \frac{\theta(\lambda \omega^n; p) \prod_{k=2}^n \theta(y_1 \omega^2 / y_k, x_k / y_1; p)}{\theta(\lambda \omega^{n-1}; p) \theta(\omega; p)^{2n-2}} Z_{n-1}^{8\text{VSOS}}(x_2, \dots, x_n; y_2, \dots, y_n; \lambda, p), \end{aligned} \quad (4.9)$$

see e.g. [R, Lemma 3.3], and the crossing symmetry

$$\begin{aligned} Z_n^{\text{8VSOS}}(\omega^2/x_1, \dots, \omega^2/x_n; 1/y_1, \dots, 1/y_n; \omega^{2n}/\lambda, p) \\ = \frac{\omega^{n(n-1)}\theta(\lambda; p)Y^n}{\theta(\lambda\omega^n; p)X^n} Z_n^{\text{8VSOS}}(x; y; \lambda, p), \end{aligned} \quad (4.10)$$

where

$$X = x_1 \cdots x_n, \quad Y = y_1 \cdots y_n.$$

The equation (4.10) can be derived from a corresponding symmetry of the Boltzmann weights, and is also apparent from the explicit formulas for the partition function given in [R].

The main result of [R] is the explicit expression

$$\begin{aligned} Z_n^{\text{8VSOS}}(x; y; \lambda, p) &= \frac{(-1)^{\binom{n}{2}}\theta(\lambda\omega^n; p)}{\theta(\omega; p)^{n^2}\theta(\gamma; p)^n Y^{n+1}\theta(X\lambda\gamma\omega^n/Y; p)} \\ &\times \frac{\prod_{i,j=1}^n y_j^2 \theta(x_i/y_j, \omega x_i/y_j; p)}{\prod_{1 \leq i < j \leq n} x_j y_j \theta(x_i/x_j, y_i/y_j; p)} \\ &\times \sum_{S \subseteq \{1, \dots, n\}} (-1)^{|S|} \frac{\theta(\lambda\gamma\omega^{n-|S|}; p)}{\theta(\lambda\omega^{n-|S|}; p)} \det_{1 \leq i, j \leq n} \left(\frac{\theta(\gamma x_i^S/y_j; p)}{\theta(x_i^S/y_j; p)} \right), \end{aligned} \quad (4.11)$$

where

$$x_i^S = \begin{cases} x_i \omega, & i \in S, \\ x_i, & i \notin S \end{cases}$$

and γ is arbitrary. This can be viewed as an analogue of the Izergin–Korepin formula for the six-vertex model. We will not work with this formula directly, though we need the following immediate consequence.

Proposition 4.3 ([R, Corollary 5.4]). *As a function of λ ,*

$$\theta(\lambda\omega^{n+1}, \lambda\omega^{n+2}; p) Z_n^{\text{8VSOS}}(x; y; \lambda, p)$$

is analytic on $\mathbb{C}^$. More precisely, it is an A_1 theta function of nome p and norm $\omega^{2n}Y/X$.*

By (4.8), it follows that

$$\frac{\theta(\lambda^3; p^3)^{n^2+2n+3}}{\theta(\lambda\omega^2, \lambda\omega^{n+1}; p)^2} Z_n^{3C} \left(\frac{1}{\theta(\lambda; p)^3}, \frac{1}{\theta(\lambda\omega; p)^3}, \frac{1}{\theta(\lambda\omega^2; p)^3} \right) \quad (4.12)$$

is analytic in $\lambda \in \mathbb{C}^*$. This remarkable fact is a key result for the present work. We do not know how to prove it except as a consequence of (4.11).

We also need the following functional equation, which was recently obtained by Razumov and Stroganov.

Proposition 4.4 ([RS]). *Let*

$$F_n(x; y; \lambda, p) = \theta(\lambda\omega^{n+1}, \lambda\omega^{n+2}; p)\Delta(x, y; p)Z_n^{\text{SVSOS}}(\omega x; y; \lambda; p),$$

where

$$\Delta(x_1, \dots, x_N; p) = \prod_{1 \leq i < j \leq N} x_j \theta(x_i/x_j; p). \quad (4.13)$$

Then,

$$\sum_{k=0}^2 F_n(\omega^k x_1, x_2, \dots, x_n; y; \omega^{-k} \lambda, p) = 0.$$

Consider F_n as a function of λ . By Proposition 4.3, it is an A_1 theta function of nome p and norm $\omega^n Y/X$. By (4.3), it can be decomposed as

$$F_n(x; y; \lambda, p) = F_n^{(0)}(x; y; p)\theta(-\omega^n \lambda^2 Y/X; p^2) + F_n^{(1)}(x; y; p)\lambda\theta(-p\omega^n \lambda^2 Y/X; p^2).$$

The following result is then immediate from Proposition 4.4.

Corollary 4.5. *For $i = 0, 1$,*

$$\sum_{k=0}^2 F_n^{(i)}(\omega^k x_1, x_2, \dots, x_n; y) = 0.$$

5. PROOF OF THEOREM 3.1

Consider (4.12) as a function of λ . By Proposition 4.3, it is an A_1 theta function of nome p and norm ω^n . Since the space of such functions is spanned by $\theta(\lambda\omega^{2n}; p)^2$ and $\theta(\lambda\omega^{2n+1}, \lambda\omega^{2n+2}; p)$, there exist functions X_n and Y_n such that

$$\begin{aligned} \frac{\theta(\lambda^3; p^3)^{n^2+2n+3}}{\theta(\lambda\omega^2, \lambda\omega^{n+1}; p)^2} Z_n^{\text{C}}(t_0, t_1, t_2) \\ = X_n(p)\theta(\lambda\omega^{2n}; p)^2 + Y_n(p)\theta(\lambda\omega^{2n+1}, \lambda\omega^{2n+2}; p), \end{aligned} \quad (5.1)$$

where t_i are as in (4.7). We will see that this decomposition corresponds to Theorem 3.1, the p -dependence being encoded in the polynomials q_n and r_n .

We first give some preliminary results.

Lemma 5.1. *There exists a function $p \mapsto \tau(p)$ such that, under the parametrization (4.7),*

$$\frac{1}{t_0} + \frac{1}{t_1} + \frac{1}{t_2} = \tau(p)\theta(\lambda^3; p^3). \quad (5.2)$$

Moreover, the quantity $T(t_0, t_1, t_2)$, defined in (3.1), equals $\tau(p)^3$.

Proof. Each of the functions $1/t_i = \theta(\lambda\omega^i; p)^3$, as well as $\theta(\lambda^3; p^3)$, are A_2 theta functions of nome p and norm 1. Thus, the first statement can be reduced to the trivial verification that $\sum_i 1/t_i$ vanishes at the points $\lambda = 1, \omega, \omega^2$. The second statement follows using

$$t_0 t_1 t_2 = \frac{1}{\theta(\lambda^3; p^3)^3}. \quad (5.3)$$

□

It was pointed out to us by Don Zagier that

$$\tau(p) = 3 \left(1 + 9p \frac{(p^9; p^9)_\infty^3}{(p; p)_\infty^3} \right), \quad (5.4)$$

where we use the notation (4.5). We also have (see the remark after Lemma 9.1)

$$\tau(p)^3 = 27 \left(1 + 27p \frac{(p^3; p^3)_\infty^{12}}{(p; p)_\infty^{12}} \right). \quad (5.5)$$

This is consistent with (5.4) in view of the identity

$$\left(1 + 9p \frac{(p^9; p^9)_\infty^3}{(p; p)_\infty^3} \right)^3 = 1 + 27p \frac{(p^3; p^3)_\infty^{12}}{(p; p)_\infty^{12}},$$

which can be found in Ramanujan's notebooks [Be, p. 345]. We mention that the function

$$p \frac{(p^3; p^3)_\infty^{12}}{(p; p)_\infty^{12}}$$

is well-known in the theory of modular forms. It is automorphic under the group

$$\Gamma_0(3) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{SL}(2, \mathbb{Z}); c \equiv 0 \pmod{3} \right\}$$

(acting by $\tau \mapsto (a\tau + b)/(c\tau + d)$, where $p = e^{2\pi i\tau}$) and in fact generates the field of all such functions [S, Thm. 21].

Lemma 5.2. *Let f be a homogeneous rational function in three variables. Suppose that, under the parametrization (4.7), $f(t_0, t_1, t_2) \equiv 0$. Then, $f \equiv 0$.*

Proof. It is enough to show that the Jacobian of the map $(\lambda, \mu, p) \mapsto (\mu t_0, \mu t_1, \mu t_2)$ does not vanish identically. Clearing denominators, the Jacobian is proportional to

$$\det \begin{pmatrix} \theta(\lambda; p) & \theta'_\lambda(\lambda; p) & \theta'_p(\lambda; p) \\ \theta(\lambda\omega; p) & \omega\theta'_\lambda(\lambda\omega; p) & \theta'_p(\lambda\omega; p) \\ \theta(\lambda\omega^2; p) & \omega^2\theta'_\lambda(\lambda\omega^2; p) & \theta'_p(\lambda\omega^2; p) \end{pmatrix}.$$

Using that

$$\theta(x; p) = (1 - x) \left(1 - (x + x^{-1})p + (1 - x - x^{-1})p^2 \right) + \mathcal{O}(p^3),$$

one can check that the determinant is

$$\frac{3\sqrt{3}i(1 + 2\lambda^3)(1 - \lambda^3)}{\lambda^2} p + \mathcal{O}(p^2).$$

□

Lemma 5.3. *Let f be a Laurent polynomial in three variables, homogeneous of degree 0. Suppose that, under the parametrization (4.7), $f(t_0, t_1, t_2)$ is independent of λ . Then, f is a polynomial in T .*

Proof. By the change of variables $\lambda \mapsto \omega\lambda$, it is seen that $f(t_0, t_1, t_2) = f(t_1, t_2, t_0)$. A priori, this holds when t_i are as in (4.7), but by Lemma 5.2 it is valid in general. Similarly, $\lambda \mapsto \lambda^{-1}$ gives $f(t_0, t_1, t_2) = f(t_0, t_2, t_1)$. Since these two transformations generate S_3 , f is symmetric, and can thus be expressed as

$$f(t_0, t_1, t_2) = \sum_{k+2l+3m=0, k, l \geq 0} C_{klm} (t_0 + t_1 + t_2)^k (t_0 t_1 + t_0 t_2 + t_1 t_2)^l (t_0 t_1 t_2)^m.$$

Introducing the function

$$\phi(\lambda, p) = (t_0 + t_1 + t_2) \theta(\lambda^3; p^3)^3$$

and using (5.2) and (5.3) gives

$$f(t_0, t_1, t_2) = \sum_{k \geq 0} \frac{\phi(\lambda, p)^k}{\theta(\lambda^3; p^3)^k} \sum_{2l+3m=-k, l \geq 0} C_{klm} \tau(p)^l.$$

We observe that, since

$$\phi(1, p) = \theta(\omega, \omega^2; p)^3 \neq 0,$$

the k th term has a pole at $\lambda = 1$ of order exactly k . In particular, if f is independent of λ , then all terms with $k \neq 0$ vanish. Thus,

$$f(t_0, t_1, t_2) = \sum_{2l+3m=0, l \geq 0} C_{0lm} (t_0 t_1 + t_0 t_2 + t_1 t_2)^l (t_0 t_1 t_2)^m,$$

which is indeed a polynomial in T . $\square$

We are now ready to prove Theorem 3.1. Assume that $n \equiv 0 \pmod{3}$. Then, (5.1) can be written

$$\begin{aligned} Z_n^{3C}(t_0, t_1, t_2) &= \frac{1}{\theta(\lambda^3; p^3)^{n(n+2)}} \left(\frac{X_n(p)}{\theta(\lambda^3; p^3)} + \frac{Y_n(p)}{\theta(\lambda; p)^3} \right) \\ &= (t_0 t_1 t_2)^{\frac{n(n+2)}{3}} \left(\frac{X_n(p) \tau(p) t_0 t_1 t_2}{t_0 t_1 + t_0 t_2 + t_1 t_2} + Y_n(p) t_0 \right), \end{aligned}$$

where $\tau(p)$ is as in Lemma 5.1. This implies

$$X_n(p) \tau(p) = \frac{t_0 t_1 + t_0 t_2 + t_1 t_2}{(t_0 t_1 t_2)^{\frac{n(n+2)}{3} + 1}} \cdot \frac{t_1 Z_n(t_0, t_1, t_2) - t_0 Z_n(t_1, t_0, t_2)}{t_1 - t_0}. \quad (5.6)$$

Since the right-hand side vanishes when $T = 0$, it follows from Lemma 5.3 that $X_n(p) \tau(p)$ is a polynomial in T divisible by T . Similarly,

$$Y_n(p) = \frac{1}{(t_0 t_1 t_2)^{\frac{n(n+2)}{3}}} \cdot \frac{Z_n(t_0, t_1, t_2) - Z_n(t_1, t_0, t_2)}{t_1 - t_0},$$

so $Y_n(p)$ is a polynomial in T . Writing

$$X_n(p) \tau(p) = (-1)^{n+1} T q_n(T), \quad (5.7a)$$

$$Y_n(p) = (-1)^n 2^{\chi(n \text{ odd})} r_n(T), \quad (5.7b)$$

we obtain Theorem 3.1 for $n \equiv 0 \pmod{3}$. The cases $n \equiv \pm 1 \pmod{3}$ are similar and we do not give the details. For later use we note that in both cases

$$X_n(p) = (-1)^{n+1} q_n(T), \quad (5.7c)$$

$$Y_n(p) \tau(p)^2 = (-1)^n 2^{\chi(n \text{ odd})} T r_n(T). \quad (5.7d)$$

Finally, we comment on the claim made before stating Proposition 3.4, that q_n , r_n and $2r_{2n+1}$ have integer coefficients. Consider the case of q_n when $n \equiv 0 \pmod{3}$; all other cases follow similarly. By (5.6) and (5.7a),

$$\begin{aligned} & \frac{t_1 Z_n(t_0, t_1, t_2) - t_0 Z_n(t_1, t_0, t_2)}{t_1 - t_0} \\ &= (-1)^{n+1} (t_0 t_1 t_2)^{\frac{n(n+2)}{3}-1} (t_0 t_1 + t_0 t_2 + t_1 t_2)^2 q_n \left(\frac{(t_0 t_1 + t_0 t_2 + t_1 t_2)^3}{(t_0 t_1 t_2)^2} \right). \end{aligned}$$

The left-hand side is a symmetric polynomial in t_i with integer coefficients. By [M2, I.2.4], it can be expanded as an integer linear combination of elementary symmetric polynomials. By the identity above, the non-zero coefficients in that expansion are coefficients of the polynomial q_n .

6. THE FUNCTION Φ_n

In §5, we expressed the function (4.12) in terms of the basis

$$(\theta(\lambda \omega^{2n}; p)^2, \theta(\lambda \omega^{2n+1}, \lambda \omega^{2n+2}; p)).$$

The first main idea for analyzing the partition function further is suggested by the trigonometric case $p = 0$. The relevant space is then the polynomials $A + B\lambda + C\lambda^2$ such that $C = \omega^n A$. From (2.5), it appears that the most natural basis to use is not $((1 - \lambda \omega^{2n})^2, (1 - \lambda \omega^{2n+1})(1 - \lambda \omega^{2n+2}))$ but rather $(1 + \omega^n \lambda^2, \lambda)$. An elliptic analogue of the latter basis is $(\theta(-\omega^n \lambda^2; p^2), \lambda \theta(-p \omega^n \lambda^2; p^2))$. As we will eventually see, this change of basis corresponds to expressing Z_n^{3C} in terms of the polynomials p_n and $\tilde{p}_n$ rather than q_n and r_n .

The next main idea is to consider a one-parameter extension of Z_n^{3C} , which is given by $Z_n^{8\text{VSOS}}$ with λ , p and x_1 free but all other parameters fixed. This corresponds to incorporating $p_n(\zeta)$ in the two-variable polynomial $P_n(x, \zeta)$.

Combining these two ideas, we consider the function

$$\theta(\lambda \omega^{n+1}, \lambda \omega^{n+2}; p) Z_n^{8\text{VSOS}}(\omega t, \omega, \dots, \omega; 1, \dots, 1; \lambda, p). \quad (6.1)$$

By Proposition 4.3 and (4.3), it is a linear combination of $\theta(-\omega^n \lambda^2/t; p^2)$ and $\lambda \theta(-p \omega^n \lambda^2/t; p^2)$, with coefficients independent of λ . It will be convenient to write the corresponding decomposition as

$$\begin{aligned} Z_n^{8\text{VSOS}}(\omega t, \omega, \dots, \omega; 1, \dots, 1; \lambda, p) &= \frac{1}{\theta(\lambda \omega^{n+1}, \lambda \omega^{n+2}; p) \theta(t; p)^{2n-1}} \\ &\times \left(p^{\frac{3n-4}{2}} t^{\frac{9n-4}{2}} \theta(-\omega^n \lambda^2/t; p^2) \tilde{\Phi}_n(t) - p^{-1} t^{\frac{3n-2}{2}} \omega^{2n} \lambda \theta(-p \omega^n \lambda^2/t; p^2) \Phi_n(t) \right). \end{aligned}$$

for n even, while for odd n we write

$$Z_n^{\text{SVSOS}}(\omega t, \omega, \dots, \omega; 1, \dots, 1; \lambda, p) = \frac{1}{\theta(\lambda\omega^{n+1}, \lambda\omega^{n+2}; p)\theta(t; p)^{2n-1}} \\ \times \left(t^{\frac{3n-1}{2}} \theta(-\omega^n \lambda^2/t; p^2) \Phi_n(t) + p^{\frac{3n-3}{2}} t^{\frac{9n-5}{2}} \omega^{2n} \lambda \theta(-p\omega^n \lambda^2/t; p^2) \tilde{\Phi}_n(t) \right).$$

The functions Φ_n and $\tilde{\Phi}_n$ depend implicitly on p but are independent of λ .

Lemma 6.1. *The functions Φ_n and $\tilde{\Phi}_n$ are related by $\tilde{\Phi}_n(t) = \Phi_n(pt)$. Moreover,*

$$\Phi_n(p^2 t) = \frac{1}{p^{6n-4} t^{6n-3}} \Phi_n(t). \quad (6.2)$$

Proof. This follows easily from the fact that, as a function of t , (6.1) is an A_{n-1} theta function of nome p and norm ω^n/λ , see [R, Lemma 3.2]. $\square$

As we will see, Φ_n is uniquely determined by the following properties.

Proposition 6.2. *The function Φ_n has the following properties:*

- (i) Φ_n is a BC_{3n-2} theta function of nome p^2 ;
- (ii) $\sum_{k=0}^2 \omega^k \Phi_n(\omega^k t) = 0$, that is, $\Phi_n(t) = f(t^3) + tg(t^3)$, with f and g analytic on $\mathbb{C}^*$;
- (iii) $t = 1$ and $t = p$ are zeroes of $\Phi_n(t)$ of multiplicity at least $2n - 1$;
- (iv) $\Phi_n(\omega) = -p^{1+3[\frac{n-2}{2}]} \omega \theta(\omega; p)^{2n-1} \lim_{t \rightarrow p} \frac{\Phi_{n-1}(t)}{\theta(t; p)^{2n-3}}$;
- (v) $\Phi_1(t) = \frac{\omega \theta(t, \pm pt; p^2)}{\theta(p; p^2)}$.

Proof. We have already noted the quasi-periodicity (6.2). For (i), it remains to show that $\Phi_n(t^{-1}) = -t^{-1} \Phi_n(t)$. This is a special case of (4.10).

Property (ii) is a special case of Corollary 4.5, where we should note that

$$\Phi_n(t) = \begin{cases} t^{(3n-2)/2} \frac{F_n^{(0)}(x; y)}{\Delta(\hat{x}, y; p)} \Big|_{x_1=pt, x_2=\dots=x_n=y_1=\dots=y_n=1}, & n \text{ even,} \\ t^{(1-3n)/2} \frac{F_n^{(0)}(x; y)}{\Delta(\hat{x}, y; p)} \Big|_{x_1=t, x_2=\dots=x_n=y_1=\dots=y_n=1}, & n \text{ odd,} \end{cases}$$

where Δ is as in (4.13) and the hats indicate omission of x_1 .

Property (iii) is obvious, and property (iv) is a special case of (4.9).

Finally, to check (v), we note that

$$F_1(x; y; \lambda) = y \theta(\omega^2 \lambda, x/y; p) R_{-+}^+(\lambda, \omega x/y) = \omega x \theta(x/y, \omega^2 \lambda, \omega^2 \lambda y/x; p).$$

By (4.3), this gives

$$F_1^{(0)}(x; y) = \omega x \theta(x/y; p) \frac{\theta(-px/y; p^2)}{\theta(p; p^2)},$$

$$\Phi_1(t) = \frac{1}{t} F_1^{(0)}(t; 1) = \frac{\omega \theta(t; p) \theta(-pt; p^2)}{\theta(p; p^2)} = \frac{\omega \theta(t, \pm pt; p^2)}{\theta(p; p^2)}.$$

□

Finally, we use (4.8) to express Z_n^{3C} in terms of Φ_n .

Corollary 6.3. *When t_i are as in (4.7) and n is even, then*

$$Z_n^{3C}(t_0, t_1, t_2) = -\omega^{n(n+1)} \frac{\theta(\lambda \omega^2, \lambda \omega^{n+1}; p)^2}{\theta(\lambda^3; p^3)^{n^2+2n+3}}$$

$$\times \left(p^{\frac{3n-4}{2}} \theta(-\omega^n \lambda^2; p^2) \lim_{t \rightarrow p} \frac{\Phi_n(t)}{\theta(t; p)^{2n-1}} + p^{-1} \omega^{2n} \lambda \theta(-p \omega^n \lambda^2; p^2) \lim_{t \rightarrow 1} \frac{\Phi_n(t)}{\theta(t; p)^{2n-1}} \right),$$

while if n is odd

$$Z_n^{3C}(t_0, t_1, t_2) = \omega^{n(n+1)} \frac{\theta(\lambda \omega^2, \lambda \omega^{n+1}; p)^2}{\theta(\lambda^3; p^3)^{n^2+2n+3}}$$

$$\times \left(\theta(-\omega^n \lambda^2; p^2) \lim_{t \rightarrow 1} \frac{\Phi_n(t)}{\theta(t; p)^{2n-1}} - p^{\frac{3n-3}{2}} \omega^{2n} \lambda \theta(-p \omega^n \lambda^2; p^2) \lim_{t \rightarrow p} \frac{\Phi_n(t)}{\theta(t; p)^{2n-1}} \right).$$

7. SYMMETRIC FUNCTIONS

We are now faced with two problems: to construct a function satisfying all properties of Proposition 6.2, and to show that this function is unique. The key for solving both problems is to replace property (iii) by a generic vanishing condition. This leads to certain symmetric multivariable theta functions, which after uniformization become the symmetric polynomials (3.9).

7.1. Symmetric theta functions. We denote by V_n the space of BC_{3n+1} theta functions of nome p^2 satisfying property (ii) of Proposition 6.2. For $t_i \in \mathbb{C}^*/\{t = p^2 t = t^{-1}\}$, we denote by $V_n(t_1, \dots, t_k)$ the subspace of V_n consisting of functions ψ such that, apart from the trivial zeroes at 1 and $\pm p$, ψ vanishes at $t_1, \dots, t_k$ (counted with multiplicity). Then, properties (i)–(iii) of Proposition 6.2 can be summarized as

$$\Phi_n \in V_{n-1}(\underbrace{1, \dots, 1}_{n-1}, \underbrace{p, \dots, p}_{n-1}). \quad (7.1)$$

Lemma 7.1. *The space V_n has dimension $2n + 1$. The space $V_n(t_1, \dots, t_k)$ has dimension at least $2n + 1 - k$, with equality for $k \leq n$.*

Proof. A basis for the BC_{3n+1} theta functions of nome p^2 is given by

$$t^{j-3n-1}\theta(-p^{2j}t^{6n+3}; p^{12n+6}) - t^{3n+2-j}\theta(-p^{2j}t^{-6n-3}; p^{12n+6}), \quad 1 \leq j \leq 3n+1,$$

see [RSc, Proposition 6.1]. It is clear that the subspace V_n is spanned by the $2n+1$ basis vectors with $j \not\equiv 0 \pmod{3}$.

Since $V_n(t_1, \dots, t_k)$ is obtained by imposing k linear conditions on V_n , it has dimension at least $2n+1-k$. For the final statement, note that by the quintuple product identity (4.4), V_n contains all functions of the form

$$\theta(t, \pm pt; p^2)\theta(b_1 t^{\pm 3}, \dots, b_n t^{\pm 3}; p^6).$$

When $b_i = t_i$ for $1 \leq i \leq k$ and the remaining b_i are generic, this function is in $V_n(t_1, \dots, t_k) \setminus V_n(t_1, \dots, t_{k+1})$. Thus, as long as we impose at most n vanishing conditions, each additional condition decreases the dimension by one. $\square$

Although we do not need it in full generality, the reader may find the following characterization of the space V_n helpful.

Lemma 7.2. *Fix $\alpha \in \mathbb{C}^*/\{t = p^2 t = t^{-1}\}$ with $\alpha \neq -\omega$. Then, V_n is the space of all functions ψ that can be written*

$$\begin{aligned} \psi(t) = \theta(t, \pm pt; p^2) \big(A \theta(-t^{\pm}, -\omega t^{\pm}, \alpha t^{\pm}; p^2) \theta(a_1 t^{\pm 3}, \dots, a_{n-1} t^{\pm 3}; p^6) \\ + B \theta(b_1 t^{\pm 3}, \dots, b_n t^{\pm 3}; p^6) \big), \end{aligned}$$

with $A, B \in \mathbb{C}$ and $a_1, \dots, a_{n-1}, b_1, \dots, b_n \in \mathbb{C}^*$.

Proof. It is easy to check that any BC_{3n+1} theta function ψ of nome p^2 can be written

$$\psi(t) = \theta(t, \pm pt; p^2) (f(t^3) + tg(t^3) + t^{-1}g(t^{-3})),$$

where f is a D_{n+1} theta function of nome p^6 , and g satisfies

$$g(p^6 t) = \frac{1}{p^{6n+2} t^{2n}} g(t). \quad (7.2)$$

Since the term involving f has the desired form, we restrict to the case $f \equiv 0$. By the quintuple product identity (4.4), $\psi \in V_n$ if and only if

$$\theta(-p^2 t; p^6) g(t^{-1}) = t \theta(-p^2 t^{-1}; p^6) g(t).$$

Together with (7.2), this implies that g vanishes on all zeroes of $\theta(-t, -p^2 t; p^6)$. Factoring $g(t) = t^{-1} \theta(-t, -p^2 t; p^6) h(t)$ gives

$$\psi(t) = t^{-2} \theta(t, \pm pt; p^2) \theta(-t^3; p^6) (\theta(-p^2 t^3; p^6) + t \theta(-p^2 t^{-3}; p^6)) h(t^3),$$

where h is a D_n theta function of nome p^6 .

Next, we observe that $\theta(-p^2 t^3; p^6) + t \theta(-p^2 t^{-3}; p^6)$ vanishes on the zeroes of $\theta(-t; p^2)$. Thus

$$\theta(-p^2 t^3; p^6) + \theta(-p^2 t^{-3}; p^6) = t \theta(-t; p^2) k(t),$$

where k is a D_2 theta function of nome p^2 , that is, $k(t) = C\theta(\beta t^\pm; p^2)$ for some C and β (depending on p). After simplification, we conclude that

$$\psi(t) = C\theta(t, \pm pt; p^2)\theta(-t^\pm, -\omega t^\pm, \beta t^\pm; p^2)h(t^3).$$

Using (4.1) to write

$$\theta(\beta t^\pm; p^2) = A\theta(\alpha t^\pm; p^2) + B\theta(-\omega^2 t^\pm; p^2),$$

we arrive at an expression of the desired form. This shows that any $\psi \in V_n$ can be expressed as indicated. The converse follows by similar arguments. $\square$

In view of (7.1), we are mainly interested in the space $V_n(t_1, \dots, t_{2n})$. Generically, one expects it to be one-dimensional and spanned by the alternant

$$\psi(t) = \det_{1 \leq i, j \leq 2n+1} (\psi_j(t_i)),$$

with $(\psi_j)_{j=1}^{2n+1}$ a basis of V_n and $t_{2n+1} = t$. However, we have not found such expressions useful for our purposes. Instead, we will work with the following less symmetric determinants.

Theorem 7.3. *Fixing a basis ψ_1, ψ_2, ψ_3 of V_1 , define*

$$\Psi_1(t_1, t_2, t_3) = \det_{1 \leq i, j \leq 3} (\psi_j(t_i))$$

and, more generally,

$$\Psi_n(t_1, \dots, t_n, u_1, \dots, u_n, v) = \frac{\prod_{i,j=1}^n u_j^{-3} \theta(u_j^3 t_i^{\pm 3}; p^6)}{\theta(v, \pm pv; p^2)^{n-1}} \det_{1 \leq i, j \leq n} \left(\frac{\Psi_1(t_i, u_j, v)}{u_j^{-3} \theta(u_j^3 t_i^{\pm 3}; p^6)} \right).$$

Let $(\psi_j^{(n)})_{j=1}^{2n+1}$ be a basis of V_n . Then,

$$\Psi_n(t_1, \dots, t_{2n+1}) = C \det_{1 \leq i, j \leq 2n+1} (\psi_j^{(n)}(t_i)),$$

with C independent of each t_i . In particular, Ψ_n is anti-symmetric in all $2n+1$ variables.

Note that Ψ_n is only defined up to a multiplicative constant, which we do not specify.

Remark 7.4. If we choose the basis $\psi_j^{(n)}$ as in the proof of Lemma 7.1, then the alternant is a minor of the determinant corresponding to the BC_{3n+1} Macdonald identity. This should mean that Ψ_n can be interpreted as an affine Lie algebra character. Although this observation may have interesting consequences, we will not explore it here. In §7.4, we will see that the trigonometric limit case $p = 0$ corresponds to characters of the orthogonal and symplectic groups. The corresponding limit of Theorem 7.3 is closely related to some determinant identities of Okada [O], see (7.11).

We divide the proof of Theorem 7.3 into a few lemmas.

Lemma 7.5. *For any $i = 1, \dots, 2n$, the map $t_i \mapsto \Psi_n(t_1, \dots, t_{2n+1})$ is an element of $V_n(t_1, \dots, \hat{t}_i, \dots, t_{2n+1})$.*

The proof of Lemma 7.5 is straight-forward. It will follow from Theorem 7.3 that the statement holds also for $i = 2n + 1$.

Lemma 7.6. *If the map $t \mapsto \Psi_n(t, t_1, \dots, t_{2n})$ is not identically zero, then*

$$\dim V_n(t_1, \dots, t_{2n}) = 1.$$

Proof. Take t_0 with $\Psi_n(t_0, t_1, \dots, t_{2n}) \neq 0$. Consider the functions

$$\psi_k(t) = \Psi_n(t_0, t_1, \dots, t_{k-1}, t, t_{k+1}, \dots, t_{2n}), \quad k = 0, 1, \dots, 2n - 1.$$

By Lemma 7.5, these functions all belong to the space $V_n(t_{2n})$. Moreover, they are linearly independent since

$$\psi_k(t_j) \neq 0 \iff j = k, \quad j, k = 0, \dots, 2n - 1.$$

Since, by Lemma 7.1, $\dim V_n(t_{2n}) = 2n$, we conclude that $\psi_0, \dots, \psi_{2n-1}$ span $V_n(t_{2n})$.

Suppose now that $f \in V_n(t_1, \dots, t_{2n})$. In particular, $f \in V_n(t_{2n})$, so we can expand

$$f(t) = \sum_{k=0}^{2n-1} A_k \psi_k(t).$$

The remaining vanishing conditions for f give $A_1 = \dots = A_{2n-1} = 0$. Thus, f is proportional to ψ_0 , so $\dim V_n(t_1, \dots, t_{2n}) = 1$. $\square$

Proof of Theorem 7.3. Let

$$C(t_1, \dots, t_{2n+1}) = \frac{\Psi_n(t_1, \dots, t_{2n+1})}{\det_{1 \leq i, j \leq 2n+1} (\psi_j^{(n)}(t_i))}.$$

We first prove that C is independent of t_1 . By Lemma 7.5, the denominator and numerator are both in $V_n(t_2, \dots, t_{2n+1})$. If that space is one-dimensional, C is independent of t_1 . Otherwise, Lemma 7.6 gives $C \equiv 0$. The same argument applies to the variables $t_2, \dots, t_{2n}$. Thus,

$$\Psi_n(t_1, \dots, t_{2n+1}) = C(t_{2n+1}) \det_{1 \leq i, j \leq 2n+1} (\psi_j^{(n)}(t_i)),$$

where it remains to show that C is independent of t_{2n+1} .

To complete the proof, we write

$$\frac{C(v)}{C(t_1)} = - \frac{\Psi_n(t_1, \dots, t_n, u_1, \dots, u_n, v)}{\Psi_n(v, t_2, \dots, t_n, u_1, \dots, u_n, t_1)} \quad (7.3)$$

and specialize $u_n = \omega t_n$. It follows from the definition of Ψ_n that

$$\begin{aligned} \Psi_n(t_1, \dots, t_n, u_1, \dots, u_{n-1}, \omega t_n, v) &= \frac{\prod_{j=1}^n t_n^{-3} u_j^{-3} \theta(t_n^3 t_j^{\pm 3}, u_j^3 t_n^{\pm 3}; p^6)}{\theta(v, \pm p v; p^2)} \\ &\times \Psi_1(t_n, \omega t_n, v) \Psi_{n-1}(t_1, \dots, t_{n-1}, u_1, \dots, u_{n-1}, v). \end{aligned}$$

Using this in (7.3) gives

$$\begin{aligned} \frac{C(v)}{C(t_1)} &= -\frac{\theta(t_1, \pm pt_1; p^2)\theta(t_n^3 t_1^{\pm 3}; p^6)\Psi_1(t_n, \omega t_n, v)}{\theta(v, \pm pv; p^2)\theta(t_n^3 v^{\pm 3}; p^6)\Psi_1(t_n, \omega t_n, t_1)} \\ &\quad \times \frac{\Psi_{n-1}(t_1, \dots, t_{n-1}, u_1, \dots, u_{n-1}, v)}{\Psi_{n-1}(v, t_2, \dots, t_{n-1}, u_1, \dots, u_{n-1}, t_1)}. \end{aligned}$$

We claim that

$$\frac{\theta(t_1, \pm pt_1; p^2)\theta(t_n^3 t_1^{\pm 3}; p^6)\Psi_1(t_n, \omega t_n, v)}{\theta(v, \pm pv; p^2)\theta(t_n^3 v^{\pm 3}; p^6)\Psi_1(t_n, \omega t_n, t_1)} = 1.$$

To see this, consider the denominator and numerator as functions of v . They both belong to the space $V_1(t_n, \omega t_n)$, which is one-dimensional for generic t_n . Thus, the quotient is independent of v and can be computed by letting $v = t_1$.

We have now reduced (7.3) to

$$\frac{C(v)}{C(t_1)} = -\frac{\Psi_{n-1}(t_1, \dots, t_{n-1}, u_1, \dots, u_{n-1}, v)}{\Psi_{n-1}(v, t_2, \dots, t_{n-1}, u_1, \dots, u_{n-1}, t_1)}.$$

By iteration, we conclude that

$$\frac{C(v)}{C(t_1)} = -\frac{\Psi_1(t_1, u_1, v)}{\Psi_1(v, u_1, t_1)} = 1.$$

This completes the proof of Theorem 7.3. $\square$

7.2. Uniformization. The next step is uniformization, see §4.1. We will work with the uniformizing map

$$\xi(t) = \frac{\theta(-p\omega; p^2)^2 \theta(\omega t^\pm; p^2)}{\theta(-\omega; p^2)^2 \theta(p\omega t^\pm; p^2)} \quad (7.4)$$

and write

$$\zeta = \frac{\omega^2 \theta(-1, -p\omega; p^2)}{\theta(-p, -\omega; p^2)}. \quad (7.5)$$

Lemma 7.7. *We have*

$$\begin{aligned} \xi(-1) &= 1, & \xi(-\omega) &= \zeta, \\ \xi(1) &= 2\zeta + 1, & \xi(p) &= \frac{\zeta}{\zeta + 2}. \end{aligned}$$

Proof. The first two identities are obvious. By (4.6),

$$\xi(s) - \xi(t) = -\frac{\omega \theta(p, p\omega; p^2) \theta(-p\omega; p^2)^2 \theta(st^\pm; p^2)}{s \theta(-\omega; p^2)^2 \theta(p\omega s^\pm, p\omega t^\pm; p^2)}. \quad (7.6)$$

Plugging in $s = 1$ and $t = -1$, and using (4.2), we obtain $\xi(1) - 1 = 2\zeta$. The case $s = p$ and $t = -1$ similarly gives $\xi(p) - 1 = -2\xi(p)/\zeta$. $\square$

We denote by W_n the space of polynomials q of degree at most $3n$ such that

$$\psi(t) = \theta(t, \pm pt; p^2) \theta(p\omega t^\pm; p^2)^{3n} q(\xi(t)),$$

is an element of V_n . Then, the correspondence between ψ and q is a bijection. We denote by $W_n(x_1, \dots, x_k)$ the subspace of W_n such that $\psi \in V_n(t_1, \dots, t_k)$, where $x_i = \xi(t_i)$.

Lemma 7.8. *The space W_1 is spanned by the three polynomials f , g and h defined in (3.6).*

Proof. By Lemma 7.2, V_1 is spanned by the functions

$$\theta(p^3 t^{\pm 3}; p^6), \quad \theta(-t^\pm, -\omega t^\pm, p\omega t^\pm; p^2), \quad \theta(t^{\pm 3}; p^6).$$

By Lemma 7.7, the uniformization of these functions are indeed, up to multiplicative constants, respectively

$$(\zeta + 2)x - \zeta, \quad (x - 1)(x - \zeta), \quad x^2(x - (2\zeta + 1)).$$

□

Taking determinants of the basis elements (3.6), we get the following results.

Corollary 7.9. *For any elements q_1, q_2, q_3 of W_1 ,*

$$\det_{1 \leq i, j \leq 3} (q_j(x_i)) = C(x_2 - x_1)(x_3 - x_1)(x_3 - x_2)F(x_1, x_2, x_3),$$

where C is a constant and F is as in (3.7).

Moreover, writing $x = \xi(t)$ and $y = \xi(u)$,

$$u^{-3} \theta(u^3 t^{\pm 3}; p^6) \theta(p\omega t^\pm, p\omega u^\pm; p^2) = C(y - x)G(x, y),$$

where C is a constant and G is as in (3.8).

It follows that the function Ψ_n from Theorem 7.3 is given by

$$\begin{aligned} \Psi_n(t_1, \dots, t_{2n+1}) &= C \prod_{i=1}^{2n+1} \theta(t_i, \pm pt_i; p^2) \theta(p\omega t_i^\pm; p^2)^n \prod_{1 \leq i < j \leq 2n+1} t_j^{-1} \theta(t_j t_i^\pm; p^2) \\ &\quad \times S_n(x_1, \dots, x_{2n+1}), \end{aligned} \quad (7.7)$$

where C is a constant, $x_i = \xi(t_i)$, and S_n is the polynomial introduced in (3.9). In particular, S_n is symmetric in all variables.

Finally, we note the following symmetry of S_n .

Lemma 7.10. *Indicating also the dependence on ζ , the polynomial S_n satisfies*

$$S_n(x_1^{-1}, \dots, x_{2n+1}^{-1}; \zeta^{-1}) = \frac{1}{\zeta^{2n^2} x_1^n \dots x_{2n+1}^n} S_n(x_1, \dots, x_{2n+1}; \zeta)$$

Proof. This follows from an elementary computation, using

$$F(x^{-1}, y^{-1}, z^{-1}; \zeta^{-1}) = \frac{1}{\zeta^2 xyz} F(x, y, z; \zeta),$$

$$G(x^{-1}, y^{-1}; \zeta^{-1}) = \frac{1}{\zeta^2 x^2 y^2} G(x, y; \zeta).$$

□

7.3. Recursions. Note that any minor of the determinant in (3.9) is a determinant of the same type. Thus, any algebraic relation between minors (see e.g. [N, Chapter 6]) implies a relation involving the polynomials S_n . It is not our purpose to give an exhaustive list of such identities; we only mention a few examples that will be used below or otherwise seem of particular interest.

First, we apply Jacobi's identity (sometimes called the Lewis Carroll formula after one of its proponents)

$$X_{n-1,n}^{n-1,n} X = X_n^n X_{n-1}^{n-1} - X_{n-1}^n X_n^{n-1},$$

where $X = \det_{1 \leq i, j \leq n} (x_{ij})$ and the other quantities are minors, upper and lower indices signifying omitted rows and columns. Choosing

$$x_{ij} = \frac{F(x_i, y_j, z)}{G(x_i, y_j)},$$

one obtains after relabelling the following recursion for the polynomials S_n .

Lemma 7.11. *For $\mathbf{x} = (x_1, \dots, x_{2n-1})$,*

$$\begin{aligned} (a-b)(c-d)S_{n-1}(\mathbf{x})S_{n+1}(a, b, c, d, \mathbf{x}) \\ = G(a, d)G(b, c)S_n(a, c, \mathbf{x})S_n(b, d, \mathbf{x}) - G(a, c)G(b, d)S_n(a, d, \mathbf{x})S_n(b, c, \mathbf{x}). \end{aligned}$$

One of the Plücker relations is

$$X_{n,n+1}^n X_{n-1} - X_{n-1,n+1}^n X_n + X_{n-1,n}^n X_{n+1} = 0,$$

valid for minors of an $n \times (n+1)$ matrix. Specializing the matrix entries as above yields the following result.

Lemma 7.12. *For $\mathbf{x} = (x_1, \dots, x_{2n-2})$,*

$$\begin{aligned} (c-d)G(a, b)S_{n-1}(b, \mathbf{x})S_n(a, c, d, \mathbf{x}) + (d-b)G(a, c)S_{n-1}(c, \mathbf{x})S_n(a, b, d, \mathbf{x}) \\ + (b-c)G(a, d)S_{n-1}(d, \mathbf{x})S_n(a, b, c, \mathbf{x}) = 0. \end{aligned}$$

Combining Lemmas 7.11 and 7.12, we obtain the following recursion.

Corollary 7.13. *For $\mathbf{x} = (x_1, \dots, x_{2n-2})$,*

$$\begin{aligned} & (b-d)(y-a)(y-c)S_{n-1}(a, \mathbf{x})S_{n-1}(c, \mathbf{x})S_{n+1}(y, a, b, c, d, \mathbf{x}) \\ & + (a-y)G(b, c)G(y, d)S_{n-1}(c, \mathbf{x})S_n(a, c, d, \mathbf{x})S_n(y, a, b, \mathbf{x}) \\ & + (y-c)G(a, d)G(y, b)S_{n-1}(a, \mathbf{x})S_n(a, b, c, \mathbf{x})S_n(y, c, d, \mathbf{x}) \\ & + (c-a)G(y, b)G(y, d)S_n(a, b, c, \mathbf{x})S_n(a, c, d, \mathbf{x})S_{n-1}(y, \mathbf{x}) = 0. \end{aligned}$$

Proof. The sum of the first two terms can be written

$$\begin{aligned} & (y-a)S_{n-1}(c, \mathbf{x})\{(b-d)(y-c)S_{n-1}(a, \mathbf{x})S_{n+1}(y, a, b, c, d, \mathbf{x}) \\ & - G(b, c)G(y, d)S_n(a, c, d, \mathbf{x})S_n(y, a, b, \mathbf{x})\}. \end{aligned}$$

By Lemma 7.11, with $(\mathbf{x}, a, b, c, d) \mapsto ((\mathbf{x}, a), d, b, c, y)$, the factor in brackets equals

$$-G(c, d)G(y, b)S_n(a, b, c, \mathbf{x})S_n(y, a, d, \mathbf{x}).$$

The sum of the last two terms is

$$\begin{aligned} & G(y, b)S_n(a, b, c, \mathbf{x})\{(y-c)G(a, d)S_{n-1}(a, \mathbf{x})S_n(y, c, d, \mathbf{x}) \\ & + (c-a)G(y, d)S_n(a, c, d, \mathbf{x})S_{n-1}(y, \mathbf{x})\}. \end{aligned}$$

By Lemma 7.12, with $(\mathbf{x}, a, b, c, d) \mapsto (\mathbf{x}, d, a, y, c)$, the factor in brackets is

$$(y-a)G(c, d)S_{n-1}(c, \mathbf{x})S_n(y, a, d, \mathbf{x}).$$

It is now clear that the sum of all four terms vanishes. $\square$

Corollary 7.13 is particularly interesting in the case when

$$\mathbf{x} = (\underbrace{a, \dots, a}_{n-1}, \underbrace{b, \dots, b}_{n-1}),$$

$c = b$ and $d = a$. As we will see, one special case is the three-term recursion for the polynomials P_n .

Corollary 7.14. *For fixed a and b , the polynomials*

$$S_n(y) = S_n(y, \underbrace{a, \dots, a}_n, \underbrace{b, \dots, b}_n)$$

satisfy

$$\begin{aligned} & (b-a)(y-a)(y-b)S_{n-1}(a)S_{n-1}(b)S_{n+1}(y) \\ & + \left((a-y)G(b, b)G(y, a)S_{n-1}(b)S_n(a) \right. \\ & \quad \left. + (y-b)G(a, a)G(y, b)S_{n-1}(a)S_n(b) \right) S_n(y) \\ & + (b-a)G(y, a)G(y, b)S_n(a)S_n(b)S_{n-1}(y) = 0. \end{aligned}$$

We also consider a similar specialization of Lemma 7.11.

Corollary 7.15. *For fixed a, b , and z , the polynomials*

$$S_n(x, y) = S_n(x, y, z, \underbrace{a, \dots, a}_{n-1}, \underbrace{b, \dots, b}_{n-1})$$

satisfy

$$\begin{aligned} (x-a)(y-b)S_{n-1}(a, b)S_{n+1}(x, y) \\ = G(a, y)G(b, x)S_n(a, b)S_n(x, y) - G(a, b)G(x, y)S_n(x, b)S_n(a, y). \end{aligned}$$

Two cases of Corollary 7.15 are of special interest. The first one is $x = b$ and $y = a$, when it reads

$$\begin{aligned} (a-b)^2 S_{n-1}(a, b)S_{n+1}(a, b) \\ = G(a, b)^2 S_n(a, a)S_n(b, b) - G(a, a)G(b, b)S_n(a, b)^2. \end{aligned} \quad (7.8)$$

This will yield Theorem 3.7. The second one is the limit case $x \rightarrow a, y \rightarrow b$, where we first divide through with $(x-a)(y-b)$ and then use l'Hôpital's rule on the right-hand side. Let

$$T_n(x, y) = S_n(\underbrace{x, \dots, x}_n, \underbrace{y, \dots, y}_n, z).$$

Since

$$\left. \frac{\partial T_n}{\partial x} \right|_{x=a, y=b} = n \left. \frac{\partial S_n}{\partial x} \right|_{x=a, y=b}$$

and similarly for y , the result can be expressed in terms of T_n and its derivatives.

Corollary 7.16. *The polynomials $T_n = T_n(x, y)$ satisfy*

$$T_{n-1}T_{n+1} = \left(\frac{\partial G}{\partial x} \frac{\partial G}{\partial y} - G \frac{\partial^2 G}{\partial x \partial y} \right) T_n^2 + \frac{1}{n^2} G^2 \left(T_n \frac{\partial^2 T_n}{\partial x \partial y} - \frac{\partial T_n}{\partial x} \frac{\partial T_n}{\partial y} \right).$$

Equivalently, if we let

$$\tau_n = \frac{1}{\prod_{k=0}^{n-1} (k!)^2} \cdot \frac{T_n}{G^{n^2}},$$

then τ_n satisfies the two-dimensional Toda molecule equation

$$\frac{\partial^2}{\partial x \partial y} \log \tau_n = \frac{\tau_{n-1} \tau_{n+1}}{\tau_n^2}.$$

We have recovered an instance of the well-known fact that this equation can be solved by determinants, which can be found already in Darboux's classic treatise [D, §378] (we owe this reference to Jacques Perk), see [H, LS] for more recent accounts.

The one-dimensional Toda equation

$$\frac{\partial^2}{\partial x^2} \log \tau_n = \frac{\tau_{n-1} \tau_{n+1}}{\tau_n^2}$$

plays an important role in the analysis of the six-vertex model with domain wall boundary conditions [BF, BL1, BL2, BL3, KZ, Zi]. This suggests that Corollary 7.16 might be useful for proving Conjecture 3.14.

7.4. Trigonometric limit. In the trigonometric limit $p \rightarrow 0$, (7.5) reduces to $\zeta = -2$. We will show that the corresponding limit of the function S_n is related to symplectic and odd orthogonal characters.

Let $\lambda = (\lambda_1, \dots, \lambda_n)$ be a partition. Recall [FH, §24.2] that the characters of the Lie algebras $\mathfrak{sp}(2n)$ and $\mathfrak{so}(2n+1)$ are given by

$$\chi_{\lambda}^{\mathfrak{sp}(2n)}(t_1, \dots, t_n) = \frac{\det_{1 \leq i, j \leq n} \left(t_j^{-(\lambda_i + n - i + 1)} - t_j^{\lambda_i + n - i + 1} \right)}{\prod_{i=1}^n t_i^{-n} (1 - t_i^2) \prod_{1 \leq i < j \leq n} (t_j - t_i)(1 - t_i t_j)},$$

$$\chi_{\lambda}^{\mathfrak{so}(2n+1)}(t_1, \dots, t_n) = \frac{\det_{1 \leq i, j \leq n} \left(t_j^{-(\lambda_i + n - i + \frac{1}{2})} - t_j^{\lambda_i + n - i + \frac{1}{2}} \right)}{\prod_{i=1}^n t_i^{\frac{1}{2} - n} (1 - t_i) \prod_{1 \leq i < j \leq n} (t_j - t_i)(1 - t_i t_j)}.$$

In this notation, we have the following result.

Theorem 7.17. *Let the variables t_i , u_i , x_i and y_i be related by*

$$x_i = -(1 + t_i + t_i^{-1}), \quad y_i = \frac{3}{1 + u_i + u_i^{-1}}.$$

Then,

$$\begin{aligned} \lim_{\zeta \rightarrow -2} (\zeta + 2)^{2mn - m(m-1)} S_n \left(x_1, \dots, x_{2n+1-2m}, \frac{\zeta y_1}{\zeta + 2}, \dots, \frac{\zeta y_{2m}}{\zeta + 2} \right) \\ = 2^{(2m+n)n - m(m-1)} 3^{2mn - m(m-1)} \prod_{i=1}^{2m} \frac{1}{(1 + u_i + u_i^{-1})^n} \\ \times \chi_{(n-m, n-m, n-m-1, n-m-1, \dots, 1, 1, 0)}^{\mathfrak{so}(4n-4m+3)}(t_1, \dots, t_{2n-2m+1}) \\ \times \chi_{(m-1, m-1, \dots, 1, 1, 0, 0)}^{\mathfrak{sp}(4m)}(u_1, \dots, u_{2m}), \end{aligned} \quad (7.9)$$

$$\begin{aligned} \lim_{\zeta \rightarrow -2} (\zeta + 2)^{(2m+1)n - m^2} S_n \left(x_1, \dots, x_{2n-2m}, \frac{\zeta y_1}{\zeta + 2}, \dots, \frac{\zeta y_{2m+1}}{\zeta + 2} \right) \\ = 2^{(2m+n+1)n - m^2} 3^{(2m+1)n - m^2} \prod_{i=1}^{2m+1} \frac{1}{(1 + u_i + u_i^{-1})^n} \\ \times \chi_{(n-m, n-m-1, n-m-1, \dots, 1, 1, 0)}^{\mathfrak{so}(4n-4m+1)}(t_1, \dots, t_{2n-2m}) \\ \times \chi_{(m, m-1, m-1, \dots, 1, 1, 0, 0)}^{\mathfrak{sp}(4m+2)}(u_1, \dots, u_{2m+1}). \end{aligned} \quad (7.10)$$

Note that the correspondence between x_i and t_i is the trigonometric limit of the uniformization (7.4):

$$\lim_{p \rightarrow 0} \xi(t_i) = -(1 + t_i + t_i^{-1}) = x_i.$$

For the variables y_i and u_i , we have instead

$$\left(\frac{\zeta \xi(u_i)}{\zeta + 2} \right)^{-1} \Big|_{\zeta \mapsto \zeta^{-1}} = \frac{1 + 2\zeta}{\xi(u_i)} \rightarrow \frac{3}{1 + u_i + u_i^{-1}} = y_i, \quad p \rightarrow 0,$$

which is a natural limit in view of Lemma 7.10.

We will use some determinant identities due to Okada [O]. Let

$$W^n(x_1, \dots, x_n; a_1, \dots, a_n) = \det_{1 \leq i, j \leq n} (x_i^{j-1} + a_i x_i^{n-j}).$$

(In [O], W^n denotes the matrix rather than its determinant.) Then,

$$\begin{aligned} \det_{1 \leq i, j \leq n} \left(\frac{W^2(x_i, y_j; a_i, b_j)}{(1 - x_i y_j)(y_j - x_i)} \right) &= \frac{1}{\prod_{i, j=1}^n (1 - x_i y_j)(y_j - x_i)} \\ &\times W^{2n}(x_1, \dots, x_n, y_1, \dots, y_n; a_1, \dots, a_n, b_1, \dots, b_n), \end{aligned} \quad (7.11a)$$

$$\begin{aligned} \det_{1 \leq i, j \leq n} \left(\frac{W^3(x_i, y_j, z; a_i, b_j, c)}{(1 - x_i y_j)(y_j - x_i)} \right) &= \frac{(1 + c)^{n-1}}{\prod_{i, j=1}^n (1 - x_i y_j)(y_j - x_i)} \\ &\times W^{2n+1}(x_1, \dots, x_n, y_1, \dots, y_n, z; a_1, \dots, a_n, b_1, \dots, b_n, c). \end{aligned} \quad (7.11b)$$

Okada used these identities to enumerate certain symmetry classes of alternating sign matrices.

The characters appearing in Theorem 7.17 can be obtained as special cases of Okada's determinants.

Lemma 7.18. *The following identities hold:*

$$\frac{W^{2n+1}(t_1^3, \dots, t_{2n+1}^3; -t_1, \dots, -t_{2n+1})}{\prod_{i=1}^{2n+1} t_i^n (1 - t_i) \prod_{1 \leq i < j \leq 2n+1} (t_j - t_i)(1 - t_i t_j)} = \chi_{(n, n, n-1, n-1, \dots, 1, 1, 0)}^{\mathfrak{so}(4n+3)}(t_1, \dots, t_{2n+1}),$$

$$\frac{(-1)^n W^{2n}(t_1^3, \dots, t_{2n}^3; -t_1, \dots, -t_{2n})}{\prod_{i=1}^{2n} t_i^{n-1} (1 - t_i^2) \prod_{1 \leq i < j \leq 2n} (t_j - t_i)(1 - t_i t_j)} = \chi_{(n-1, n-1, \dots, 1, 1, 0, 0)}^{\mathfrak{sp}(4n)}(t_1, \dots, t_{2n}),$$

$$\frac{W^{2n+1}(t_1^3, \dots, t_{2n+1}^3; -t_1^2, \dots, -t_{2n+1}^2)}{\prod_{i=1}^{2n+1} t_i^n (1 - t_i^2) \prod_{1 \leq i < j \leq 2n+1} (t_j - t_i)(1 - t_i t_j)} = \chi_{(n, n-1, n-1, \dots, 1, 1, 0, 0)}^{\mathfrak{sp}(4n+2)}(t_1, \dots, t_{2n+1}),$$

$$\frac{(-1)^n W^{2n}(t_1^3, \dots, t_{2n}^3; -t_1^2, \dots, -t_{2n}^2)}{\prod_{i=1}^{2n} t_i^n (1 - t_i) \prod_{1 \leq i < j \leq 2n} (t_j - t_i)(1 - t_i t_j)} = \chi_{(n, n-1, n-1, \dots, 1, 1, 0)}^{\mathfrak{so}(4n+1)}(t_1, \dots, t_{2n}).$$

Proof. This is straight-forward, and we only provide some details for the first identity. By definition,

$$\begin{aligned} W^{2n+1}(t_1^3, \dots, t_{2n+1}^3; -t_1, \dots, -t_{2n+1}) &= \det_{1 \leq i, j \leq 2n+1} \left(t_i^{3(j-1)} - t_i^{3(2n+1-j)+1} \right) \\ &= \prod_{i=1}^{2n+1} t_i^{\frac{6n+1}{2}} \det_{1 \leq i, j \leq 2n+1} \left(t_i^{\mu_j} - t_i^{-\mu_j} \right), \end{aligned}$$

where $\mu_j = 3n - 3j + 7/2$. In order to make all μ_j positive, we multiply the last n rows by -1 . To sort the μ_j in descending order, each of those rows must be moved an odd number of positions upwards. Taken together, these two operations do not change the determinant, and we arrive at the desired result. $\square$

We will need the following elementary identities.

Lemma 7.19. *In the notation above, the following identities hold:*

$$\begin{aligned} x_2 - x_1 &= \frac{(1 - t_1 t_2)(t_2 - t_1)}{t_1 t_2}, \\ y_2 - y_1 &= \frac{3(1 - u_1)(1 - u_2)(1 - u_1 u_2)(u_2 - u_1)}{(1 - u_1^3)(1 - u_2^3)}, \\ \lim_{\zeta \rightarrow -2} G(x_1, x_2) &= \frac{2(t_2^3 - t_1^3)(1 - t_1^3 t_2^3)}{t_1^2 t_2^2 (t_2 - t_1)(1 - t_1 t_2)}, \\ \lim_{\zeta \rightarrow -2} (\zeta + 2)^2 G\left(x_1, \frac{\zeta y_1}{\zeta + 2}\right) &= \frac{72 u_1^2 (1 - u_1)^2}{(1 - u_1^3)^2}, \\ \lim_{\zeta \rightarrow -2} (\zeta + 2)^2 G\left(\frac{\zeta y_1}{\zeta + 2}, \frac{\zeta y_2}{\zeta + 2}\right) &= \frac{72(1 - u_1)^2 (1 - u_2)^2 (u_2^3 - u_1^3)(1 - u_1^3 u_2^3)}{(1 - u_1^3)^2 (1 - u_2^3)^2 (u_2 - u_1)(1 - u_1 u_2)}, \\ \lim_{\zeta \rightarrow -2} F(x_1, x_2, x_3) &= \frac{2 W^3(t_1^3, t_2^3, t_3^3; -t_1, -t_2, -t_3)}{\prod_{i=1}^3 t_i (1 - t_i) \prod_{1 \leq i < j \leq 3} (t_j - t_i)(1 - t_i t_j)}, \\ \lim_{\zeta \rightarrow -2} (\zeta + 2) F\left(x_1, x_2, \frac{\zeta y_1}{\zeta + 2}\right) &= -\frac{12 u_1 (1 - u_1) W^2(t_1^3, t_2^3; -t_1^2, -t_2^2)}{(1 - u_1^3) t_1 (1 - t_1) t_2 (1 - t_2) (t_2 - t_1) (1 - t_1 t_2)}, \\ \lim_{\zeta \rightarrow -2} (\zeta + 2)^2 F\left(x_1, \frac{\zeta y_1}{\zeta + 2}, \frac{\zeta y_2}{\zeta + 2}\right) &= -\frac{72 u_1 u_2 W^2(u_1^3, u_2^3; -u_1, -u_2)}{(1 + u_1)(1 - u_1^3)(1 + u_2)(1 - u_2^3)(u_2 - u_1)(1 - u_1 u_2)}, \\ \lim_{\zeta \rightarrow -2} (\zeta + 2)^2 F\left(\frac{\zeta y_1}{\zeta + 2}, \frac{\zeta y_2}{\zeta + 2}, \frac{\zeta y_3}{\zeta + 2}\right) &= \frac{72 W^3(u_1^3, u_2^3, u_3^3; -u_1^2, -u_2^2, -u_3^2)}{\prod_{i=1}^3 (1 + u_i)(1 - u_i^3) \prod_{1 \leq i < j \leq 3} (u_j - u_i)(1 - u_i u_j)}. \end{aligned}$$

Proof of Theorem 7.17. To prove (7.9), we relabel the variables, considering instead

$$\lim_{\zeta \rightarrow -2} (\zeta + 2)^{2mn - m(m-1)} \times S_n \left(\frac{\zeta x_1}{\zeta + 2}, \dots, \frac{\zeta x_m}{\zeta + 2}, x_{m+1}, \dots, x_n, \frac{\zeta y_1}{\zeta + 2}, \dots, \frac{\zeta y_m}{\zeta + 2}, y_{m+1}, \dots, y_n, z \right). \quad (7.12)$$

Applying (3.9) leads to the block determinant

$$\det \begin{pmatrix} \frac{F\left(\frac{\zeta x_i}{\zeta+2}, \frac{\zeta y_j}{\zeta+2}, z\right)}{G\left(\frac{\zeta x_i}{\zeta+2}, \frac{\zeta y_j}{\zeta+2}\right)} & \frac{F\left(\frac{\zeta x_i}{\zeta+2}, y_j, z\right)}{G\left(\frac{\zeta x_i}{\zeta+2}, y_j\right)} \\ \frac{F\left(x_i, \frac{\zeta y_j}{\zeta+2}, z\right)}{G\left(x_i, \frac{\zeta y_j}{\zeta+2}\right)} & \frac{F(x_i, y_j, z)}{G(x_i, y_j)} \end{pmatrix}.$$

By Lemma 7.19, the off-diagonal blocks vanish in the limit $\zeta \rightarrow -2$, so the determinant splits as the product of the diagonal blocks. It follows that (7.12) can be written

$$\begin{aligned} \lim_{\zeta \rightarrow -2} \prod_{\substack{1 \leq i \leq m, \\ m+1 \leq j \leq n}} \frac{(\zeta + 2)^4 G\left(x_i, \frac{\zeta y_j}{\zeta+2}\right) G\left(\frac{\zeta x_i}{\zeta+2}, y_j\right)}{(\zeta + 2)^2 \left(x_j - \frac{\zeta x_i}{\zeta+2}\right) \left(y_j - \frac{\zeta y_i}{\zeta+2}\right)} \\ \times (\zeta + 2)^{m(m+1)} S_m \left(\frac{\zeta x_1}{\zeta + 2}, \dots, \frac{\zeta x_m}{\zeta + 2}, \frac{\zeta y_1}{\zeta + 2}, \dots, \frac{\zeta y_m}{\zeta + 2}, z \right) \\ \times S_{n-m}(x_{m+1}, \dots, x_n, y_{m+1}, \dots, y_n, z). \end{aligned}$$

Computing the prefactor using Lemma 7.19, we are reduced to proving the two cases $m = 0$ and $m = n$ of (7.9), that is,

$$\lim_{\zeta \rightarrow -2} S_n(x_1, \dots, x_{2n+1}) = 2^{n^2} \chi_{(n,n,n-1,n-1,\dots,1,1,0)}^{\mathfrak{so}(4n+3)}(t_1, \dots, t_{2n+1}),$$

$$\begin{aligned} \lim_{\zeta \rightarrow -2} (\zeta + 2)^{n(n+1)} S_n \left(\frac{\zeta y_1}{\zeta + 2}, \dots, \frac{\zeta y_{2n}}{\zeta + 2}, z \right) \\ = 2^{n(2n+1)} 3^{n(n+1)} \prod_{i=1}^{2n} \frac{1}{(1 + u_i + u_i^{-1})^n} \chi_{(n-1,n-1,\dots,1,1,0,0)}^{\mathfrak{sp}(4n)}(u_1, \dots, u_{2n}). \end{aligned}$$

This is now straight-forward, using Lemma 7.19, Okada's identities (7.11) and finally Lemma 7.18.

The proof of (7.10) is similar. We consider the limit

$$\lim_{\zeta \rightarrow -2} (\zeta + 2)^{(2m+1)n-m^2} \times S_n \left(\frac{\zeta x_1}{\zeta + 2}, \dots, \frac{\zeta x_m}{\zeta + 2}, x_{m+1}, \dots, x_n, \frac{\zeta y_1}{\zeta + 2}, \dots, \frac{\zeta y_m}{\zeta + 2}, y_{m+1}, \dots, y_n, \frac{\zeta z}{\zeta + 2} \right).$$

Incorporating a factor $(\zeta + 2)^m$ into the determinant, we consider

$$\lim_{\zeta \rightarrow -2} \det \begin{pmatrix} \frac{(\zeta + 2)F\left(\frac{\zeta x_i}{\zeta + 2}, \frac{\zeta y_j}{\zeta + 2}, \frac{\zeta z}{\zeta + 2}\right)}{G\left(\frac{\zeta x_i}{\zeta + 2}, \frac{\zeta y_j}{\zeta + 2}\right)} & \frac{(\zeta + 2)F\left(\frac{\zeta x_i}{\zeta + 2}, y_j, \frac{\zeta z}{\zeta + 2}\right)}{G\left(\frac{\zeta x_i}{\zeta + 2}, y_j\right)} \\ \frac{F\left(x_i, \frac{\zeta y_j}{\zeta + 2}, \frac{\zeta z}{\zeta + 2}\right)}{G\left(x_i, \frac{\zeta y_j}{\zeta + 2}\right)} & \frac{F\left(x_i, y_j, \frac{\zeta z}{\zeta + 2}\right)}{G\left(x_i, y_j\right)} \end{pmatrix}.$$

In the limit, the upper right block vanishes. Similarly as before, this reduces the proof of (7.10) to a straight-forward verification of the special cases

$$\begin{aligned} \lim_{\zeta \rightarrow -2} (\zeta + 2)^n S_n \left(x_1, \dots, x_{2n}, \frac{\zeta y_1}{\zeta + 2} \right) \\ = \frac{2^{n(n+1)} 3^n}{(1 + u_1 + u_1^{-1})^n} \chi_{(n, n-1, n-1, \dots, 1, 1, 0)}^{\mathfrak{so}(4n+1)}(t_1, \dots, t_{2n}), \end{aligned}$$

$$\begin{aligned} \lim_{\zeta \rightarrow -2} (\zeta + 2)^{n(n+1)} S_n \left(\frac{\zeta y_1}{\zeta + 2}, \dots, \frac{\zeta y_{2n+1}}{\zeta + 2} \right) \\ = 2^{n(2n+1)} 3^{n(n+1)} \prod_{i=1}^{2n+1} \frac{1}{(1 + u_i + u_i^{-1})^n} \chi_{(n, n-1, n-1, \dots, 1, 1, 0, 0)}^{\mathfrak{sp}(4n+2)}(u_1, \dots, u_{2n+1}). \end{aligned}$$

□

As a consequence of Theorem 7.17, we have the following identities.

Corollary 7.20. *One has*

$$\begin{aligned} \lim_{\zeta \rightarrow -2} \left(1 + \frac{\zeta}{2} \right)^{n^2 - \left\lceil \frac{(n-1)^2}{4} \right\rceil} S_n \left(\underbrace{2\zeta + 1, \dots, 2\zeta + 1}_{n+1}, \underbrace{\frac{\zeta}{\zeta + 2}, \dots, \frac{\zeta}{\zeta + 2}}_n \right) \\ = \begin{cases} 2^{n^2} 3^{\frac{n^2}{4}} A_{n+1}, & n \text{ even}, \\ 2^{n^2} 3^{\frac{n^2-1}{4}} C_{n+1}, & n \text{ odd}, \end{cases} \end{aligned}$$

$$\begin{aligned}
\lim_{\zeta \rightarrow -2} \left(1 + \frac{\zeta}{2}\right)^{n(n+1) - \left[\frac{(n-1)^2}{4}\right]} S_n \left(\underbrace{2\zeta + 1, \dots, 2\zeta + 1}_n, \underbrace{\frac{\zeta}{\zeta + 2}, \dots, \frac{\zeta}{\zeta + 2}}_{n+1} \right) \\
= \begin{cases} 2^{n^2-1} 3^{\frac{n(n-2)}{4}} C_{n+1}, & n \text{ even}, \\ 2^{n^2-1} 3^{\frac{(n-1)^2}{4}} A_{n+1}, & n \text{ odd}, \end{cases}
\end{aligned}$$

where A_n and C_n are as in (1.1) and (2.6).

To verify these identities, we note that the limits correspond to special cases of Theorem 7.17 where $t_i = u_i = 1$ for all i . Recall that, if $\mu_i = \lambda_i + n - i$, then [FH]

$$\begin{aligned}
\chi_{\lambda}^{\text{sp}(2n)}(1, \dots, 1) &= \frac{\prod_{1 \leq i < j \leq n} (\mu_i - \mu_j) \prod_{1 \leq i \leq j \leq n} (\mu_i + \mu_j + 2)}{2^n 1! 3! \dots (2n-1)!}, \\
\chi_{\lambda}^{\text{so}(2n+1)}(1, \dots, 1) &= \frac{\prod_{1 \leq i < j \leq n} (\mu_i - \mu_j) \prod_{1 \leq i \leq j \leq n} (\mu_i + \mu_j + 1)}{1! 3! \dots (2n-1)!}.
\end{aligned}$$

The relevant special cases can be simplified as

$$\begin{aligned}
\chi_{(n, n, n-1, n-1, \dots, 1, 1, 0)}^{\text{so}(4n+3)}(1, \dots, 1) &= 3^{n^2} \prod_{k=0}^n \frac{(6k+1)!}{(2n+2k+1)!}, \\
\chi_{(n-1, n-1, \dots, 1, 1, 0, 0)}^{\text{sp}(4n)}(1, \dots, 1) &= 3^{n(n-1)} \prod_{k=0}^{n-1} \frac{(6k+4)!}{(2n+2k+2)!}, \\
\chi_{(n, n-1, n-1, \dots, 1, 1, 0, 0)}^{\text{sp}(4n+2)}(1, \dots, 1) &= \frac{3^{n^2}}{2^{2n+1}} \prod_{k=0}^n \frac{(6k+2)(6k)!}{(2n+2k+1)!}, \\
\chi_{(n, n-1, n-1, \dots, 1, 1, 0)}^{\text{so}(4n+1)}(1, \dots, 1) &= 2^{2n} 3^{n(n-1)} \prod_{k=0}^{n-1} \frac{(6k+5)(6k+3)!}{(2n+2k+2)!}.
\end{aligned}$$

Using these identities, it is straight-forward to derive Corollary 7.20 from Theorem 7.17.

8. THE POLYNOMIALS P_n AND p_n

In this Section, we show how specialization of the variables in S_n lead to the polynomials P_n and p_n , and deduce a number of properties of the latter two systems.

8.1. Elementary factors. Our first task is to identify some elementary factors that appear when specializing the polynomials S_n . We will use the notation

$$\delta_n = \left\lfloor \frac{n^2}{4} \right\rfloor;$$

cf. (3.2). For later reference, we mention the identities

$$\delta_{n+1} + \delta_{n-1} - 2\delta_n = \chi(n \text{ odd}), \quad (8.1)$$

$$\left\lfloor \frac{n(n+2)}{2} \right\rfloor = n^2 - 2\delta_{n-1} = 2\delta_n + n. \quad (8.2)$$

Proposition 8.1. *There exist polynomials $P_n(x, \zeta)$, $p_n(\zeta)$ and $y_n(\zeta)$ such that*

$$\begin{aligned} \left(1 + \frac{\zeta}{2}\right)^{n^2} S_n\left(x, \underbrace{2\zeta + 1, \dots, 2\zeta + 1}_n, \underbrace{\frac{\zeta}{\zeta + 2}, \dots, \frac{\zeta}{\zeta + 2}}_n\right) \\ = (-1)^{\binom{n+1}{2}} \zeta^{n^2} (1 + \zeta)^{n^2} (1 + 2\zeta)^{\delta_{n-1}} \left(1 + \frac{\zeta}{2}\right)^{\delta_{n-1}} P_n(x, \zeta), \end{aligned} \quad (8.3a)$$

$$\begin{aligned} \left(1 + \frac{\zeta}{2}\right)^{n^2} S_n\left(\underbrace{2\zeta + 1, \dots, 2\zeta + 1}_{n+1}, \underbrace{\frac{\zeta}{\zeta + 2}, \dots, \frac{\zeta}{\zeta + 2}}_n\right) \\ = (-1)^{\binom{n+1}{2}} \zeta^{n^2} (1 + \zeta)^{n^2} (1 + 2\zeta)^{\delta_n} \left(1 + \frac{\zeta}{2}\right)^{\delta_{n-1}} p_n(\zeta), \end{aligned} \quad (8.3b)$$

and, for $n \geq 1$,

$$\begin{aligned} \left(1 + \frac{\zeta}{2}\right)^{n(n-1)} S_n\left(\underbrace{2\zeta + 1, \dots, 2\zeta + 1}_{n+2}, \underbrace{\frac{\zeta}{\zeta + 2}, \dots, \frac{\zeta}{\zeta + 2}}_{n-1}\right) \\ = (-1)^{\binom{n-1}{2}} 2^{\lfloor \frac{n+5}{2} \rfloor} \zeta^{n^2-1} (1 + \zeta)^{n^2} (1 + 2\zeta)^{\delta_{n+1}} \left(1 + \frac{\zeta}{2}\right)^{\delta_{n-2}} y_n(\zeta). \end{aligned} \quad (8.3c)$$

Applying Lemma 7.10 to (8.3a) gives, using also (8.2),

$$P_n(x, \zeta) = x^n \zeta^{\lfloor \frac{n(n+2)}{2} \rfloor} P_n(1/x, 1/\zeta). \quad (8.4)$$

Similarly, applying Lemma 7.10 to (8.3b) gives

$$\begin{aligned} \left(1 + \frac{\zeta}{2}\right)^{n(n+1)} S_n\left(\underbrace{2\zeta + 1, \dots, 2\zeta + 1}_n, \underbrace{\frac{\zeta}{\zeta + 2}, \dots, \frac{\zeta}{\zeta + 2}}_{n+1}\right) \\ = (-1)^{\binom{n+1}{2}} 2^{-\lfloor \frac{n+1}{2} \rfloor} \zeta^{n(n+1)} (1 + \zeta)^{n^2} (1 + 2\zeta)^{\delta_{n-1}} \left(1 + \frac{\zeta}{2}\right)^{\delta_n} \tilde{p}_n(\zeta), \end{aligned}$$

where

$$\tilde{p}_n(\zeta) = \zeta^{n(n+1)/2} p_n(1/\zeta).$$

We will see in Proposition 8.9 that $\deg p_n = n(n+1)/2$, so the notation agrees with (3.3). Moreover, we clearly have

$$P_n(2\zeta + 1, \zeta) = (1 + 2\zeta)^{\lfloor \frac{n}{2} \rfloor} p_n(\zeta), \quad (8.5a)$$

$$P_n\left(\frac{\zeta}{\zeta+2}, \zeta\right) = \zeta^n(\zeta+2)^{-\lceil \frac{n+1}{2} \rceil} \tilde{p}_n(\zeta). \quad (8.5b)$$

To prove Proposition 8.1, we first observe that $S_n(x_1, \dots, x_{2n+1})$ is a polynomial of degree at most n in each x_i . This is obvious for the variable z in (3.9), and thus holds in general by symmetry. It follows that the left-hand sides in (8.3) are polynomials in x and ζ . We need to show that they vanish of appropriate degree at the points $\zeta = 0, -1, -1/2$ and -2 .

We first consider the function

$$S_n(1 + \zeta x_1, \dots, 1 + \zeta x_n, \zeta y_1, \dots, \zeta y_n, z). \quad (8.6)$$

Expressing it as in (3.9), we observe that since $\zeta^2 \mid G(1 + \zeta x, \zeta y)$, the prefactor $\prod_{i,j=1}^n G(1 + \zeta x_i, y_j)$ is divisible by ζ^{2n^2} . The product

$$\prod_{1 \leq i < j \leq n} ((1 + \zeta x_j) - (1 + \zeta x_i))(\zeta y_j - \zeta y_i)$$

contributes a factor $\zeta^{n(n-1)}$ to the denominator. Finally, since $\zeta \mid F(1 + \zeta x, \zeta y, z)$, each matrix entry has a single pole at $\zeta = 0$, so the determinant has a pole of degree at most n . In total, we conclude that (8.6) is divisible by $\zeta^{2n^2 - n(n-1) - n} = \zeta^{n^2}$. This implies the same statement for the left-hand sides of (8.3a) and (8.3b).

In the case of (8.3c), we consider instead

$$S_n\left(1 + \zeta x_1, \dots, 1 + \zeta x_{n-2}, x_{n-1}, x_n, \zeta y_1, \dots, \zeta y_{n-2}, y_{n-1}, \frac{\zeta}{\zeta+2}, z\right). \quad (8.7)$$

Using that $\zeta \mid G(x, \zeta y)$ and that

$$G\left(x, \frac{\zeta}{\zeta+2}\right) = \frac{2\zeta^2(\zeta+1)^2}{(\zeta+2)^2} \quad (8.8)$$

for all x , obvious modifications of the previous argument shows that (8.7), and hence also the left-hand side of (8.3c), is divisible by ζ^{n^2-1} .

The case $\zeta = -1$ is simple. Considering

$$S_n(-1 + (\zeta+1)x_1, \dots, -1 + (\zeta+1)x_n, -1 + (\zeta+1)y_1, \dots, -1 + (\zeta+1)y_n, z), \quad (8.9)$$

and observing that

$$\begin{aligned} (\zeta+1) &\mid F(-1 + (\zeta+1)x, -1 + (\zeta+1)y, z), \\ (\zeta+1)^2 &\mid G(-1 + (\zeta+1)x, -1 + (\zeta+1)y), \end{aligned}$$

one checks that (8.9) is divisible by $(\zeta+1)^{n^2}$. This implies the corresponding statement for the three left-hand sides of (8.3).

The vanishing conditions at $\zeta = -1/2$ and $\zeta = -2$ follow from the following Lemma.

Lemma 8.2. *As a polynomial in ζ ,*

$$S_n((2\zeta + 1)x_1, \dots, (2\zeta + 1)x_k, x_{k+1}, \dots, x_{2n+1})$$

is divisible by $(2\zeta + 1)^{\delta_{k-1}}$, and

$$(\zeta + 2)^{kn} S_n\left(\frac{\zeta x_1}{\zeta + 2}, \dots, \frac{\zeta x_k}{\zeta + 2}, x_{k+1}, \dots, x_{2n+1}\right)$$

is divisible by $(\zeta + 2)^{\delta_{k-1}}$.

Proof. The second statement follows from Theorem 7.17 (note that we only need the very first step in the proof, not the formulas involving Lie algebra characters). The first statement then follows using Lemma 7.10. $\square$

8.2. Recursions. We will now specialize the recursions for S_n given in §7.3 to obtain recursions for P_n and p_n .

If we let $a = 2\zeta + 1$ and $b = \zeta/(\zeta + 2)$ in (7.8), using (8.8) and the identities

$$G(2\zeta + 1, x) = 2(\zeta + 1)^2 x^2,$$

$$2\zeta + 1 - \frac{\zeta}{\zeta + 2} = \frac{2(\zeta + 1)^2}{\zeta + 2},$$

we obtain the following result after simplification.

Proposition 8.3. *The polynomials p_n and y_n satisfy*

$$\begin{aligned} (\zeta + 1)^2 p_{n+1}(\zeta) p_{n-1}(\zeta) &= \zeta^{n+1} \tilde{p}_n(\zeta) y_n(\zeta) \\ &+ (1 + 2\zeta)^{1+\chi(n \text{ even})} \left(1 + \frac{\zeta}{2}\right)^{1+\chi(n \text{ odd})} p_n(\zeta)^2. \end{aligned} \quad (8.10)$$

Equivalently, (3.5) holds, where $Y = y_n$.

By induction, we obtain the following consequence, which is a key result in our approach.

Corollary 8.4. *The polynomial p_n satisfies $p_n(0) = 1$; in particular, it does not vanish identically.*

From this, another key result follows.

Corollary 8.5. *The space $V_n(\underbrace{1, \dots, 1}_n, \underbrace{p, \dots, p}_n)$ defined in §7.1 has dimension 1.*

Proof. By Lemma 7.6 and (7.7), it is enough to show that the polynomial

$$S_n\left(x, \underbrace{2\zeta + 1, \dots, 2\zeta + 1}_n, \underbrace{\frac{\zeta}{\zeta + 2}, \dots, \frac{\zeta}{\zeta + 2}}_n\right)$$

does not vanish identically. By (8.3b), this follows from Corollary 8.4. $\square$

We remark that Corollary 8.5 can alternatively be deduced from Theorem 7.17.

Proposition 8.6. *The polynomials $P_n(x, \zeta)$ are determined from the starting values $P_0 = 1$ and $P_1 = x + \zeta$ by the recursion*

$$A_n(x, \zeta)P_{n+1}(x, \zeta) = B_n(x, \zeta)P_n(x, \zeta) + C_n(x, \zeta)P_{n-1}(x, \zeta), \quad (8.11)$$

where

$$\begin{aligned} A_n(x, \zeta) &= (1 + 2\zeta)^{\chi(n \text{ even})} \left(1 + \frac{\zeta}{2}\right)^{\chi(n \text{ even})} (x - 2\zeta - 1) \left(x - \frac{\zeta}{\zeta + 2}\right) \\ &\quad \times P_{n-1}(2\zeta + 1, \zeta) P_{n-1}\left(\frac{\zeta}{\zeta + 2}, \zeta\right), \\ B_n(x, \zeta) &= \left(1 + \frac{\zeta}{2}\right) \left(x^2(x - 2\zeta - 1) P_{n-1}\left(\frac{\zeta}{\zeta + 2}, \zeta\right) P_n(2\zeta + 1, \zeta) \right. \\ &\quad \left. - (2\zeta + 1)^2 \left(x - \frac{\zeta}{\zeta + 2}\right) P_{n-1}(2\zeta + 1, \zeta) P_n\left(\frac{\zeta}{\zeta + 2}, \zeta\right)\right), \\ C_n(x, \zeta) &= (\zeta + 1)^2 x^2 P_n(2\zeta + 1, \zeta) P_n\left(\frac{\zeta}{\zeta + 2}, \zeta\right). \end{aligned}$$

Proof. Letting $a = 2\zeta + 1$ and $b = \zeta/(\zeta + 2)$ in Corollary 7.14 gives (8.11) after simplification. The polynomials are determined by this recursion, provided that $A_n(x, \zeta)$ does not vanish identically. By (8.5), this follows from Corollary 8.4. $\square$

At this point, a remark on our approach is in order. It is easy to see that if a system of polynomials satisfies (8.11), then it can be used to construct (via uniformization) functions Φ_n satisfying the properties of Proposition 6.2. The reader may ask why we do not take (8.11) as our starting point, thus avoiding introducing the multivariable polynomials S_n . The problem is that, to use (8.11) as a definition, we must know that $A_n(x, \zeta)$ does not vanish identically. Moreover, we need to know that Φ_n is uniquely determined by the properties of Proposition 6.2. Both these facts have been deduced from Corollary 8.4. If one could find a way to deduce inductively from (8.11) that $P_n(2\zeta + 1, \zeta)$ and $P_n(\zeta/(\zeta + 2), \zeta)$ never vanish identically, then the proofs of our main results could be significantly shortened.

8.3. Further properties. Using Proposition 8.6, we can obtain quite detailed information on the polynomials P_n .

Proposition 8.7. *The polynomial $P_n(x, \zeta)$ can be written*

$$P_n(x, \zeta) = \sum_{k=0}^n f_k^n(\zeta) \zeta^{n-k} x^k, \quad (8.12)$$

where f_k^n is a polynomial of degree $2\delta_n$, with leading coefficient

$$\frac{1}{2^k} \binom{n+k}{n}. \quad (8.13)$$

In the notation (3.3), these polynomials satisfy

$$f_k^n(\zeta) = \tilde{f}_{n-k}^n(\zeta). \quad (8.14)$$

For $k > [(n+1)/2]$, $f_k^n(\zeta)$ is divisible by $(\zeta+2)^{k-[\frac{n+1}{2}]}$. Moreover,

$$f_n^n(\zeta) = \left(1 + \frac{\zeta}{2}\right)^{[\frac{n}{2}]} p_{n-1}(\zeta), \quad (8.15a)$$

$$f_{n-1}^n(\zeta) = \frac{1}{2} \left(1 + \frac{\zeta}{2}\right)^{[\frac{n}{2}]-1} (\zeta + n + 1) p_{n-1}(\zeta). \quad (8.15b)$$

Further results follow by applying (8.14) to the other statements. In particular, from (8.15a) we obtain

$$P_n(0, \zeta) = \zeta^n \left(\frac{2\zeta+1}{2}\right)^{[\frac{n}{2}]} \tilde{p}_{n-1}(\zeta). \quad (8.16)$$

Proof. From the proof of Proposition 8.1, and also from Proposition 8.6, it is clear that $P_n(x, \zeta)$ is of degree at most n in x , so we can write it as in (8.12), where a priori $f_k^n(\zeta)\zeta^{n-k}$ is a polynomial.

To show that f_k^n is a polynomial, we consider the function

$$F_n(x) = \lim_{\zeta \rightarrow 0} \frac{P_n(\zeta x, \zeta)}{\zeta^n} = \sum_{k=0}^n \lim_{\zeta \rightarrow 0} f_k^n(\zeta) x^k.$$

We will prove by induction that the limit exists and equals

$$F_n(x) = \sum_{k=0}^n \binom{n+k}{k} \frac{x^{n-k}}{2^k} = x^n {}_2F_1 \left(\begin{matrix} -n, n+1 \\ -n \end{matrix}; \frac{1}{2x} \right), \quad (8.17)$$

in standard hypergeometric notation. To this end, we divide (8.11) by ζ^{2n} and then let $\zeta \rightarrow 0$. We need that, by (8.5a) and Corollary 8.4,

$$P_n(1, 0) = p_n(0) = 1,$$

and we also write

$$\lim_{\zeta \rightarrow 0} \frac{P_n(\zeta/(\zeta+2), \zeta)}{\zeta^n} = F_n(1/2).$$

Then, the resulting identity simplifies to

$$\begin{aligned} & \left(x - \frac{1}{2}\right) F_{n-1}(1/2) F_{n+1}(x) \\ &= \left(x^2 F_{n-1}(1/2) + \left(x - \frac{1}{2}\right) F_n(1/2)\right) F_n(x) - x^2 F_n(1/2) F_{n-1}(x). \end{aligned}$$

We must show that this recursion is solved by (8.17). Equivalently, since

$$F_n(1/2) = \frac{1}{2^n} {}_2F_1 \left(\begin{matrix} -n, n+1 \\ -n \end{matrix}; 1 \right) = \frac{1}{2^{n+1}} \binom{2n+2}{n+1},$$

we need to check that

$$\left(x - \frac{1}{2}\right) F_{n+1}(x) = \left(x^2 + \frac{2n+1}{n+1} \left(x - \frac{1}{2}\right)\right) F_n(x) - \frac{2n+1}{n+1} x^2 F_{n-1}(x),$$

which can be done by a straight-forward computation.

The equation (8.17) shows that f_k^n is a polynomial, with

$$f_{n-k}^n(0) = \frac{1}{2^k} \binom{n+k}{n}. \quad (8.18)$$

It follows from (8.4), using also (8.2), that

$$f_k^n(\zeta) = \zeta^{2\delta_n} f_{n-k}^n(1/\zeta).$$

Together with (8.18), this proves that f_k^n has degree $2\delta_n$, as well as (8.13) and (8.14).

The statement on divisibility by powers of $\zeta + 2$ is equivalent to saying that

$$\lim_{\zeta \rightarrow -2} (\zeta + 2)^{\lfloor \frac{n+1}{2} \rfloor} P_n \left(\frac{\zeta x}{\zeta + 2}, \zeta \right)$$

exists finitely. This is a consequence of (8.3a) and Theorem 7.17.

To prove (8.15), we pick out the coefficients of x^{n+3} and x^{n+2} on both sides of (8.11). On the right, neither C_n nor the second of the two terms in B_n contribute. Using (8.5) and simplifying, we find that

$$\begin{aligned} p_{n-1}(\zeta) \left(x^2 - \left(\frac{\zeta}{\zeta + 2} + 2\zeta + 1 \right) x \right) (f_{n+1}^{n+1}(\zeta) x^{n+1} + \zeta f_n^{n+1}(\zeta) x^n) \\ = \left(1 + \frac{\zeta}{2} \right)^{\chi(n \text{ odd})} p_n(\zeta) x^2 (x - 2\zeta - 1) (f_n^n(\zeta) x^n + \zeta f_{n-1}^n(\zeta) x^{n-1}) + \dots, \end{aligned}$$

where the ellipsis denotes terms of order at most $n+1$ in x . Picking out the top coefficient gives the recursion

$$p_{n-1}(\zeta) f_{n+1}^{n+1}(\zeta) = \left(1 + \frac{\zeta}{2} \right)^{\chi(n \text{ odd})} p_n(\zeta) f_n^n(\zeta),$$

which is solved by (8.15a). Picking out the next coefficient then yields

$$p_{n-1}(\zeta) f_n^{n+1}(\zeta) = \left(1 + \frac{\zeta}{2} \right)^{\chi(n \text{ odd})} p_n(\zeta) f_{n-1}^n(\zeta) + \frac{1}{2} \left(1 + \frac{\zeta}{2} \right)^{\lfloor \frac{n-1}{2} \rfloor} p_{n-1}(\zeta) p_n(\zeta),$$

which is solved by (8.15b). $\square$

For the formulation of the next result, it is convenient to introduce the polynomials

$$\begin{aligned} \phi_n(x) &= {}_2F_1 \left(\begin{matrix} -n/2, -(n-1)/2 \\ n+3/2 \end{matrix}; x \right), \\ \psi_n(x) &= {}_3F_2 \left(\begin{matrix} -n/2, -(n+1)/2, (n+5)/4 \\ n+3/2, (n+1)/4 \end{matrix}; x \right) \end{aligned}$$

of degree $[n/2]$, $[(n+1)/2]$, respectively.

We will need the identities

$$(3x+1)\psi_n(x) = \frac{3(3n+1)(3n+4)}{(2n+1)(2n+3)} x\phi_{n+1}(x) + (x-1)^2\phi_{n-1}(x), \quad (8.19)$$

$$(3x+1)^2\phi_n(x) = \frac{3(3n+2)(3n+5)}{(2n+1)(2n+3)} x\psi_{n+1}(x) + (x-1)^2\psi_{n-1}(x), \quad (8.20)$$

$$\frac{3(3n+2)(3n+4)}{(2n+1)(2n+3)} x\phi_{n+1}(x) = (9x-1)\phi_n(x) + (x-1)^2\phi_{n-1}(x), \quad (8.21)$$

which are straight-forward to verify.

Proposition 8.8. *For special values of ζ , $P_n(x, \zeta)$ may be expressed as*

$$P_n(x, 0) = x^n, \quad (8.22)$$

$$P_n(x, -1) = (-1)^{\chi(n \equiv 2 \pmod{4})} 2^{\delta_{n-1}} (x-1)^n, \quad (8.23)$$

$$P_n(x, 1) = 2^{\delta_{n-1}-n} 3^{\lfloor \frac{n}{2} \rfloor} A_{n+1} (1+x)^n \phi_n \left(\frac{(3x-1)(x-3)}{3(1+x)^2} \right), \quad (8.24)$$

$$P_n(x, -2) = \begin{cases} (-3/4)^{\frac{n}{2}} A_{n+1} (1-x)^{\frac{n}{2}} \phi_n \left(\frac{x+3}{3(1-x)} \right), & n \text{ even}, \\ -2^{-n-1} 3^{\frac{n-1}{2}} C_{n+1} (1-x)^{\frac{n+1}{2}} \psi_n \left(\frac{x+3}{3(1-x)} \right), & n \text{ odd}. \end{cases} \quad (8.25)$$

Applying (8.4) to (8.25) one may also evaluate $P_n(x, -1/2)$.

Proof. The identity (8.22) follows from (8.12) and (8.18).

If we let $\zeta = -1$ in (8.11), we obtain after simplification

$$(-1)^{\chi(n \text{ even})} 2^{\chi(n \text{ odd})} P_{n-1}(-1, -1) P_{n+1}(x, -1) = (x-1) P_n(-1, -1) P_n(x, -1).$$

It is easy to check that this is solved by (8.23), using (8.1) to simplify the exponent of 2.

Similarly, if we let $\zeta = -1$ in (8.11) and substitute (8.24), we are reduced to the recursion (8.21). For this computation, it is useful to note that

$$\frac{A_n A_{n+2}}{A_{n+1}^2} = \frac{3(3n+2)(3n+4)}{4(2n+1)(2n+3)}$$

and that

$$y = \frac{(3x-1)(x-3)}{3(1+x)^2} \implies y-1 = -\frac{16x}{3(x+1)^2}, \quad 9y-1 = \frac{8(x^2-4x+1)}{(x+1)^2}.$$

To prove (8.25), we multiply (8.11) by $(1+\zeta/2)^{\lfloor \frac{n+1}{2} \rfloor}$ and then let $\zeta \rightarrow -2$. This gives

$$\begin{aligned} & (-3)^{\chi(n \text{ even})} (x+3) X_{n-1} P_{n-1}(-3, -2) P_{n+1}(x, -2) \\ &= X_n \left(-9 P_{n-1}(-3, -2) P_n(x, -2) + x^2 P_n(-3, -2) P_{n-1}(x, -2) \right), \end{aligned} \quad (8.26)$$

where

$$X_n = \lim_{\zeta \rightarrow -2} \left(1 + \frac{\zeta}{2}\right)^{\lfloor \frac{n+1}{2} \rfloor} P_n \left(\frac{\zeta}{\zeta + 2}, \zeta \right).$$

It follows from (8.3a) and Corollary 7.20 that

$$X_n = \begin{cases} (-1)^{\frac{n}{2}} 2^{-1} C_{n+1}, & n \text{ even}, \\ -2^{-1} A_{n+1}, & n \text{ odd}. \end{cases} \quad (8.27)$$

It is now straight-forward to check that substituting (8.25) in (8.26) yields (8.19) when n is odd and (8.20) when n is even. For this computation, one needs the identities

$$\begin{aligned} \frac{A_{n+2} C_n}{A_{n+1} C_{n+1}} &= \frac{3(3n+1)(3n+4)}{4(2n+1)(2n+3)}, \\ \frac{A_n C_{n+2}}{A_{n+1} C_{n+1}} &= \frac{3(3n+2)(3n+5)}{4(2n+1)(2n+3)}, \end{aligned}$$

and also that,

$$z = \frac{x+3}{3(1-x)} \implies z-1 = \frac{4x}{3(1-x)}, \quad 3z+1 = \frac{4}{1-x}.$$

□

Proposition 8.9. *The polynomial p_n has degree $n(n+1)/2$ and leading coefficient $2^{-\lfloor \frac{n+2}{2} \rfloor} \binom{2n+2}{n+1}$. Moreover, p_n assumes the special values*

$$p_n(-1) = (-1)^{\chi(n \equiv 1 \pmod{4})} 2^{\delta_{n+1}} \quad (8.28)$$

$$p_n(1) = 2^{\delta_{n+1}} A_{n+1} \quad (8.29)$$

$$p_n(-2) = \begin{cases} A_{n+1}, & n \text{ even}, \\ (-1)^{\frac{n+1}{2}} C_{n+1}, & n \text{ odd}, \end{cases} \quad (8.30)$$

$$p_n(-1/2) = \begin{cases} 2^{-\frac{1}{2}(n^2+2n+2)} C_{n+1}, & n \text{ even}, \\ (-1)^{\frac{n+1}{2}} 2^{-\frac{1}{2}(n+1)^2} A_{n+1}, & n \text{ odd}. \end{cases} \quad (8.31)$$

Proof. The first statement follows (8.13) and (8.15a). The evaluations (8.28), (8.29) and (8.30) follow from Proposition 8.8, using (8.5a). Finally, by (8.5b), (8.31) is equivalent to (8.27). □

Writing $p_n(\zeta) = \sum_{k=0}^{n(n+1)/2} a_k \zeta^k$, we have simple formulas for a_0 (Corollary 8.4) and $a_{n(n+1)/2}$. It is easy to deduce from Proposition 8.3 that

$$a_1 = \begin{cases} \frac{7n(n+2)}{8}, & n \text{ even}, \\ \frac{7(n^2+2n+3)}{8}, & n \text{ odd}. \end{cases}$$

It also seems that

$$a_{\frac{n(n+1)}{2}-1} = \begin{cases} \frac{n^2(7n+10)}{2^{(n+8)/2}(n+2)} \binom{2n+2}{n+1}, & n \text{ even,} \\ \frac{(n+1)(7n^2+3n-6)}{2^{(n+7)/2}(n+2)} \binom{2n+2}{n+1}, & n \text{ odd.} \end{cases}$$

One can probably prove this using Proposition 8.6, though we have not worked out the details. It does not seem that the other coefficients admit such simple expressions.

Finally, we use the first part of Proposition 8.9 to pick out the leading term on both sides of (8.10). This allows us to compute the leading term in the polynomials y_n . In particular, we may conclude that $\deg Y = \deg A$ in (3.5).

Corollary 8.10. *The polynomial y_n has degree $\frac{n(n-1)}{2} + 2$ and leading coefficient*

$$\frac{(2n+2)!(2n)!}{2^{n+\chi(n \text{ odd})}(n+2)!(n+1)!2n!}.$$

8.4. Zeroes. In this Section we prove Propositions 3.12 and 3.13.

Proof of Proposition 3.12. We proceed by induction on n , treating the cases of odd and even n separately. Assume that the statement holds for the zeroes of p_{2n} and p_{2n+1} . Let $a_1, \dots, a_n$ denote the real zeroes of p_{2n} , $b_1, \dots, b_{n+1}$ the real zeroes of $\tilde{p}_{2n+1}$, and $c_1, \dots, c_{n+1}$ the real zeroes of p_{2n+2} . We assume that

$$b_1 < a_1 < b_2 < a_2 < \dots < a_n < b_{n+1} < -2, \quad (8.32)$$

and need to prove

$$b_1 < c_1 < b_2 < c_2 < \dots < b_{n+1} < c_{n+1} < -2.$$

Since, by Proposition 8.9, p_{2n} has positive leading coefficient, it follows from (8.32) that $(-1)^{n+i+1}p_{2n}(b_i) > 0$. Moreover, by Proposition 8.3, if $\tilde{p}_{2n+1}(\zeta) = 0$, then

$$(\zeta+1)^2 p_{2n}(\zeta) p_{2n+2}(\zeta) = (1+2\zeta) \left(1 + \frac{\zeta}{2}\right)^2 p_{2n+1}(\zeta)^2.$$

We conclude that $(-1)^{n+i}p_{2n+2}(b_i) > 0$. Thus, p_{2n+2} has one zero between each consecutive pair of points b_i, b_{i+1} . Since, by (8.30), $p_{2n+2}(-2) > 0$, the remaining zero lies between b_{n+1} and -2 . The induction step for odd n is similar, and we do not give the details. $\square$

Lemma 8.11. *Fix ζ in the interval $-2 < \zeta < -1/2$ and assume Conjecture 3.11. Then, $p_n(\zeta)$ is negative if $n \equiv 1 \pmod{4}$ and positive else. Further, $\tilde{p}_n(\zeta)$ is negative if $n \equiv 2 \pmod{4}$ and positive else. Moreover, $P_n(0, \zeta)$ is positive if $n \equiv 0 \pmod{4}$ and negative else. Finally, the leading coefficient of $P_n(x, \zeta)$ is negative if $n \equiv 2 \pmod{4}$ and positive else.*

Proof. Since we assume that p_n does not vanish in the interval $-2 < \zeta < -1/2$, $p_n(\zeta)$ has the same sign as $p_n(-1)$, so the first statement is obtained from (8.28). The sign of $\tilde{p}_n(\zeta)$ is determined as a consequence. The remaining statements follow using (8.15a) and (8.16). $\square$

Proof of Proposition 3.13. For fixed ζ , with $-2 < \zeta < -1/2$ and $\zeta \neq -1$, let $a_1, \dots, a_{n-1}$ denote the zeroes of P_{n-1} , $b_1, \dots, b_n$ the zeroes of P_n , and $c_1, \dots, c_{n+1}$ the zeroes of P_{n+1} . By induction, we assume

$$0 < b_1 < a_1 < b_2 < a_2 < \dots < a_{n-1} < b_n, \quad (8.33)$$

and prove

$$0 < c_1 < b_1 < c_2 < \dots < b_n < c_{n+1}.$$

Using Lemma 8.11, we deduce from (8.33) that

$$(-1)^{i+\chi(n \equiv 1 \pmod{4})} P_{n-1}(b_i, \zeta) > 0.$$

It follows from (8.11) that, if $P_n(x, \zeta) = 0$, then

$$P_{n+1}(x, \zeta) = \frac{2^{\chi(n \text{ even})} \zeta (\zeta + 1)^2 x^2 \tilde{p}_n(\zeta) p_n(\zeta)}{(x - 2\zeta - 1)((\zeta + 2)x - \zeta) \tilde{p}_{n-1}(\zeta) p_{n-1}(\zeta)} P_{n-1}(x, \zeta).$$

Again using Lemma 8.11,

$$(-1)^n \frac{\tilde{p}_n(\zeta) p_n(\zeta)}{\tilde{p}_{n-1}(\zeta) p_{n-1}(\zeta)} > 0.$$

Combining these facts we find that

$$(-1)^{\chi(n \not\equiv 3 \pmod{4})+i} P_{n+1}(b_i, \zeta) > 0.$$

Thus, P_{n+1} has a zero between any two consecutive b_i . Moreover, yet again using Lemma 8.11, $P_{n+1}(0, \zeta) P_{n+1}(b_1, \zeta) < 0$, so there is an additional zero between 0 and b_1 . Finally, $P_{n+1}(b_n, \zeta)$ and the leading coefficient of $P_{n+1}(x, \zeta)$ have opposite signs, so the final zero is to the right of b_n . $\square$

9. RETURN TO THREE-COLOUR MODEL

9.1. Uniformization of Φ_n . After the long detour in §7 and §8, we are now ready to apply our results to the three-colour model. First, we express the function Φ_n , introduced in §6, in terms of the polynomials P_{n-1} . The following identities will be useful.

Lemma 9.1. *With ζ as in (7.5),*

$$\begin{aligned} 2\zeta + 1 &= \frac{\theta(-p\omega, \omega; p^2)^2}{\theta(-\omega, p\omega; p^2)^2}, \\ \zeta + 2 &= p \frac{\theta(-1, -\omega; p^2)\theta(\omega; p^2)^2}{\theta(-p, -p\omega; p^2)\theta(p\omega; p^2)^2}, \\ \zeta + 1 &= -\frac{\theta(p, -p\omega; p^2)}{\theta(-p, p\omega; p^2)}, \end{aligned} \tag{9.1}$$

$$\zeta - 1 = \frac{\theta(p, p\omega; p^2)\theta(\omega; p^2)^2}{\theta(-p, -p\omega; p^2)\theta(-\omega; p^2)^2}. \tag{9.2}$$

Proof. All four identities follow from Lemma 7.7, in the last two cases by writing $\zeta + 1 = \xi(1) - \xi(-\omega)$, $\zeta - 1 = \xi(-\omega) - \xi(-1)$ and using (7.6). $\square$

We note in passing that Lemma 9.1 can be used to prove (5.5). It is enough to check that, after elementary simplification,

$$\frac{(2\zeta + 1)(\zeta + 2)(\zeta - 1)^4}{\zeta(\zeta + 1)^4} = p\omega \frac{\theta(\omega; p^2)^{12}\theta(p\omega; p^2)^4}{\theta(-\omega, -p\omega; p^2)^8} = 3^6 p \frac{(p^3; p^3)_\infty^{12}}{(p; p)_\infty^{12}};$$

this implies (5.5) in view of (3.13).

Lemma 9.2. *One has*

$$\zeta(\zeta + 1)^4 = 2\eta^3,$$

where

$$\eta = -\frac{\theta(p; p^2)\theta(-p\omega; p^2)^2}{\theta(p\omega; p^2)\theta(-p; p^2)^2}.$$

Proof. This follows easily from (9.1), using (4.2). $\square$

Proposition 9.3. *In the notation above,*

$$\Phi_n(t) = B_n \theta(t, \pm pt; p^2) \theta(t^\pm, pt^\pm, p\omega t^\pm; p^2)^{n-1} P_{n-1}(\xi(t), \zeta), \tag{9.3}$$

where

$$B_n = \begin{cases} -\frac{p\omega^{1-n^2-n}\theta(-\omega; p^2)}{\theta(p, -p; p^2)\theta(p\omega; p^2)^{2n-2}\eta^{\frac{n^2}{4}}(2\zeta + 1)^{\frac{n-2}{2}}}, & n \text{ even}, \\ \frac{\omega^{-n^2-n}}{\theta(p; p^2)\theta(p\omega; p^2)^{2n-2}\eta^{\frac{n^2-1}{4}}(2\zeta + 1)^{\frac{n-1}{2}}}, & n \text{ odd}. \end{cases}$$

Proof. Both sides of (9.3) belong to the space

$$V_{n-1}(\underbrace{1, \dots, 1}_{n-1}, \underbrace{p, \dots, p}_{n-1}),$$

which is one-dimensional by Corollary 8.5. Thus, (9.3) holds for some constant B_n , which we compute using part (iv) of Proposition 6.2. We have

$$\Phi_n(\omega) = (-1)^{n+1} B_n \omega^{1-n} \theta(-p\omega; p^2) \theta(p; p^2)^{n-1} \theta(\omega; p^2)^{2n-1} \theta(p\omega; p^2)^{3n-2} P_{n-1}(0, \zeta),$$

$$\left. \frac{\Phi_{n-1}(t)}{\theta(t; p)^{2n-3}} \right|_{t=p} = B_{n-1} p^{2-n} \omega^{-n-1} \theta(-1; p^2) \theta(\omega; p^2)^{2n-4} P_{n-2} \left(\frac{\zeta}{\zeta+2}, \zeta \right).$$

By (8.5b) and (8.16),

$$\frac{P_{n-1}(0, \zeta)}{P_{n-2}(\zeta/(\zeta+2), \zeta)} = \zeta \left(\frac{(\zeta+2)(2\zeta+1)}{2} \right)^{\left[\frac{n-1}{2}\right]}.$$

We conclude that

$$\begin{aligned} \frac{B_n}{B_{n-1}} &= (-1)^n p^{3\left[\frac{n-2}{2}\right]+3-n} \omega^2 \frac{\theta(-1; p^2) \theta(\omega; p^2)^{2n-4}}{\theta(-p\omega; p^2) \theta(p, p\omega; p^2)^{n-1}} \frac{1}{\zeta} \left(\frac{2}{(\zeta+2)(2\zeta+1)} \right)^{\left[\frac{n-1}{2}\right]} \\ &= \begin{cases} -\frac{p\omega^{n+1}\theta(-\omega; p^2)}{\theta(-p; p^2)\theta(p\omega; p^2)^2\eta^{\frac{n}{2}}}, & n \text{ even}, \\ -\frac{p^{-1}\omega^{n-1}\theta(-p; p^2)}{\theta(-\omega; p^2)\theta(p\omega; p^2)^2\eta^{\frac{n-1}{2}}(2\zeta+1)}, & n \text{ odd}, \end{cases} \end{aligned}$$

where we used Lemma 9.1 to simplify the expression. It is clear that this recursion can be solved as indicated; the starting value $B_1 = \omega/\theta(p; p^2)$ follows from part (v) of Proposition 6.2. $\square$

Corollary 9.4. *When t_i are as in (4.7),*

$$\begin{aligned} Z_n^{3C}(t_0, t_1, t_2) &= \frac{\theta(\lambda\omega^2, \lambda\omega^{n+1}; p)^2}{\theta(p; p^2)\eta^{\frac{n^2}{4}}\theta(\lambda^3; p^3)^{n^2+2n+3}} \\ &\quad \times \left(\zeta^{\frac{n}{2}} \tilde{p}_{n-1}(\zeta) \theta(-p\omega, -\omega^n \lambda^2; p^2) - \omega^{1-n} \lambda p_{n-1}(\zeta) \theta(-\omega, -p\omega^n \lambda^2; p^2) \right) \end{aligned}$$

for n even, while for odd n

$$\begin{aligned} Z_n^{3C}(t_0, t_1, t_2) &= \frac{\theta(\lambda\omega^2, \lambda\omega^{n+1}; p)^2}{\theta(p; p^2)\eta^{\frac{n^2-1}{4}}\theta(\lambda^3; p^3)^{n^2+2n+3}} \\ &\quad \times \left(p_{n-1}(\zeta) \theta(-p, -\omega^n \lambda^2; p^2) - \omega^{-n} \lambda \zeta^{\frac{n-1}{2}} \tilde{p}_{n-1}(\zeta) \theta(-1, -p\omega^n \lambda^2; p^2) \right). \end{aligned}$$

Proof. In Corollary 6.3, express Φ_n as in Theorem 9.3. Using (8.5), we see that

$$\lim_{t \rightarrow 1} \frac{\Phi_n(t)}{\theta(t; p)^{2n-1}} = D_n p_{n-1}(\zeta),$$

$$\lim_{t \rightarrow p} \frac{\Phi_n(t)}{\theta(t; p)^{2n-1}} = E_n \tilde{p}_{n-1}(\zeta),$$

where

$$D_n = (-1)^{n+1} \theta(-p; p^2) \theta(p\omega; p^2)^{2n-2} (2\zeta+1)^{\left[\frac{n-1}{2}\right]} B_n,$$

$$E_n = p^{1-n} \omega^{1-n} \theta(-1; p^2) \theta(\omega; p^2)^{2n-2} \zeta^{n-1} (\zeta+2)^{-\left[\frac{n}{2}\right]} B_n.$$

To complete the proof, we use Lemma 9.1 to write these expressions as

$$D_n = \begin{cases} p\omega^{1-n-n^2} \frac{\theta(-\omega; p^2)}{\theta(p; p^2)\eta^{\frac{n^2}{4}}}, & n \text{ even} \\ \omega^{-n-n^2} \frac{\theta(-p; p^2)}{\theta(p; p^2)\eta^{\frac{n^2-1}{4}}}, & n \text{ odd}, \end{cases}$$

$$E_n = \begin{cases} -p^{2-\frac{3n}{2}}\omega^{-n(n+1)} \frac{\theta(-p\omega; p^2)\zeta^{\frac{n}{2}}}{\theta(p; p^2)\eta^{\frac{n^2}{4}}}, & n \text{ even} \\ p^{\frac{3}{2}(1-n)}\omega^{-n(n+1)} \frac{\theta(-1; p^2)\zeta^{\frac{n-1}{2}}}{\theta(p; p^2)\eta^{\frac{n^2-1}{4}}}, & n \text{ odd}. \end{cases}$$

□

Using (8.30) and (8.31), one may check that in the trigonometric case $p = 0$ (which implies $\eta = -1$ and $\zeta = -2$), Corollary 9.4 reduces to (2.5).

When $n = 2$, we obtain the following important consequence.

Corollary 9.5. *In the notation of Lemma 5.1,*

$$\tau(p) = \frac{\zeta^2 + 4\zeta + 1}{\eta}.$$

By Lemma 9.2, it follows that, when t_i are given by (4.7),

$$T(t_0, t_1, t_2) = \frac{2(\zeta^2 + 4\zeta + 1)^3}{\zeta(\zeta + 1)^4}. \quad (9.4)$$

Proof. Noting that

$$Z_2^{3C}(t_0, t_1, t_2) = \frac{\theta(\lambda; p)^3 + \theta(\lambda\omega^2; p)^3}{\theta(\lambda; p)^9\theta(\lambda\omega; p)^{12}\theta(\lambda\omega^2; p)^9}$$

and that

$$\tau(p) = \frac{\theta(\lambda; p)^3 + \theta(\lambda\omega; p)^3 + \theta(\lambda\omega^2; p)^3}{\theta(\lambda^3; p^3)} = \lim_{\lambda \rightarrow \omega^2} \frac{\theta(\lambda; p)^3 + \theta(\lambda\omega^2; p)^3}{\theta(\lambda^3; p^3)},$$

we express $\tau(p)$ using the case $n = 2$ of Corollary 9.4. After simplification, we obtain

$$\tau(p) = -\frac{\omega^2\theta(-p, -\omega; p^2)}{\theta(\omega; p)^2\theta(p; p^2)\eta} (\zeta^2(\zeta + 3) - (3\zeta + 1)).$$

Factoring

$$\zeta^2(\zeta + 3) - (3\zeta + 1) = (\zeta - 1)(\zeta^2 + 4\zeta + 1),$$

applying (9.2) and using (4.2b), we arrive at the stated result. □

9.2. Return to the polynomials q_n and r_n . Combining Theorem 3.1 and Corollary 9.4, one easily recovers two of our main results: Theorem 3.2 and Theorem 3.6.

As a starting point, we note the theta function identities

$$\begin{aligned} \theta(-\omega^n \lambda^2; p^2) \\ = \frac{\omega^2 \theta(-\omega; p^2)^2 \theta(\lambda \omega^{2n}; p)^2 - \omega \theta(-1; p^2) \theta(\lambda \omega^{2n+1}, \lambda \omega^{2n+2}; p)}{\theta(\omega; p)^2}, \end{aligned} \quad (9.5a)$$

$$\begin{aligned} \lambda \theta(-p \omega^n \lambda^2; p^2) \\ = \omega^{n+1} \frac{\theta(-p \omega; p^2)^2 \theta(\lambda \omega^{2n}; p)^2 - \theta(-p; p^2) \theta(\lambda \omega^{2n+1}, \lambda \omega^{2n+2}; p)}{\theta(\omega; p)^2}. \end{aligned} \quad (9.5b)$$

These can be obtained as special cases of (4.1), or simply by noting that since they relate three functions in a two-dimensional space, it is enough to verify them at the points $\lambda = \omega^n$ and $\lambda = \omega^{n+2}$.

Using (9.5) in Corollary 9.4, we obtain an expression of the form (5.1), where for n even,

$$\begin{aligned} X_n(p) &= -\frac{\omega^2 \theta(-\omega; p)}{\theta(p; p^2) \theta(\omega; p)^2 \eta^{\frac{n^2}{4}}} (p_{n-1}(\zeta) - \zeta^{\frac{n}{2}} \tilde{p}_{n-1}(\zeta)) \\ &= -\frac{p_{n-1}(\zeta) - \zeta^{\frac{n}{2}} \tilde{p}_{n-1}(\zeta)}{\eta^{\frac{n^2-4}{4}} (1 - \zeta^2)}, \\ Y_n(p) &= \frac{\omega^2 \theta(-\omega, -p; p^2)}{\theta(p; p^2) \theta(\omega; p)^2 \eta^{\frac{n^2}{4}}} \left(p_{n-1}(\zeta) - \zeta^{\frac{n+2}{2}} \tilde{p}_{n-1}(\zeta) \right) \\ &= \frac{p_{n-1}(\zeta) - \zeta^{\frac{n+2}{2}} \tilde{p}_{n-1}(\zeta)}{\eta^{\frac{n^2}{4}} (1 - \zeta)}, \end{aligned}$$

where we used Lemma 9.1 and (4.2). Similarly, when n is odd,

$$\begin{aligned} X_n(p) &= \frac{p_{n-1}(\zeta) - \zeta^{\frac{n+1}{2}} \tilde{p}_{n-1}(\zeta)}{\eta^{\frac{n^2-1}{4}} (1 - \zeta)}, \\ Y_n(p) &= -2 \frac{p_{n-1}(\zeta) - \zeta^{\frac{n-1}{2}} \tilde{p}_{n-1}(\zeta)}{\eta^{\frac{n^2-9}{4}} (1 - \zeta) (1 + \zeta)^3}. \end{aligned}$$

We now compare these expressions with (5.7). For instance, when $n \equiv 0 \pmod{6}$,

$$-X_n = \frac{p_{n-1}(\zeta) - \zeta^{\frac{n}{2}} \tilde{p}_{n-1}(\zeta)}{\eta^{\frac{n^2-4}{4}} (1 - \zeta^2)} = \frac{T q_n(T)}{\tau(p)}.$$

By Corollary 9.5 and (9.4), this can be written as

$$p_{n-1}(\zeta) - \zeta^{\frac{n}{2}} \tilde{p}_{n-1}(\zeta) = (1 - \zeta)^2 (\zeta^2 + 4\zeta + 1)^{\frac{n^2-4}{4}} \\ \times \left(\frac{\zeta(\zeta+1)^4}{2(\zeta^2 + 4\zeta + 1)^3} \right)^{\frac{n^2}{12}-1} q_n \left(\frac{2(\zeta^2 + 4\zeta + 1)^3}{\zeta(\zeta+1)^4} \right).$$

Since, by Corollary 8.4, $p_{n-1}(0) = 1$, it follows that q_n is monic of degree $n^2/12 - 1$, and that the final equation of Theorem 3.6 holds. Repeating the same argument for all cases, one obtains Theorem 3.2 and Theorem 3.6.

Finally, we comment on the deduction of Corollary 3.3 from Theorem 3.2. This is a tedious exercise, and we will only explain the case when $n \equiv 1 \pmod{6}$ and we consider squares of colour 2. In this case,

$$Z_n^{3C}(t_0, t_1, t_2) = (t_0 t_1 t_2)^{\frac{n(n+2)}{3}} \left(\frac{t_0 t_1}{t_2} q_n(T) - 2 \frac{t_0 t_1 + t_0 t_2 + t_1 t_2}{t_2} r_n(T) \right) \\ = (t_0 t_1 t_2)^{\frac{n(n+2)}{3}} \left(\frac{t_0 t_1}{t_2} T^{\frac{n^2-1}{12}} - 2 \frac{t_0 t_1 + t_0 t_2 + t_1 t_2}{t_2} T^{\frac{n^2-13}{12}} \right) + \dots,$$

the ellipsis denoting lower terms in T , hence also in t_2 . Writing

$$T = t_2 \frac{(t_0 + t_1)^3}{(t_0 t_1)^2} + \dots,$$

this can be simplified to

$$(t_0 t_1)^{\frac{n^2+4n+7}{6}} t_2^{\frac{5n^2+8n-13}{12}} (t_0 + t_1)^{\frac{n^2-9}{4}} (t_0^2 + t_1^2) + \dots.$$

Thus, the leading power of t_2 is $(5n^2 + 8n - 13)/12$, and the coefficient of

$$t_0^{\frac{n^2+4n+7}{6}+k} t_1^{\frac{5n^2+8n+11}{12}-k} t_2^{\frac{5n^2+8n-13}{12}}$$

is

$$\binom{(n^2-9)/4}{k} + \binom{(n^2-9)/4}{k-2}.$$

Repeating the same argument for each colour and each residue class of $n \pmod{6}$, one obtains Corollary 3.3.

9.3. Proof of Proposition 3.4. In this Section, we prove that r_{2n+1} has integer coefficients. We need the following fact, which is generalized in Corollary 9.8.

Lemma 9.6. *The polynomial $2^{\lceil \frac{n(n+2)}{2} \rceil} p_n(\zeta/2)$ has integer coefficients.*

For the proof, we will use a determinant formula for p_n .

Lemma 9.7. *The polynomials p_n are given by*

$$p_n(\zeta) = \frac{(-1)^{\binom{n+1}{2}} \zeta^{n^2} (\zeta + 1)^{n^2}}{2^{n^2} \prod_{j=1}^n (j-1)!^2 (1+2\zeta)^{\left[\frac{n^2}{4}\right]} \left(1 + \frac{\zeta}{2}\right)^{n^2 + \left[\frac{(n-1)^2}{4}\right]}} \times \det_{1 \leq i, j \leq n} \left(\frac{\partial^{i+j-2}}{\partial x^{i-1} \partial y^{j-1}} \Big|_{x=2\zeta+1, y=\frac{\zeta}{\zeta+2}} \frac{F(x, y, 2\zeta+1)}{G(x, y)} \right).$$

Proof. Let $x_i \rightarrow 2\zeta + 1$, $y_i \rightarrow \zeta/(\zeta + 2)$ and $z = 2\zeta + 1$ in (3.9). The left-hand side can be expressed in terms of the polynomials p_n using (8.3b). We now apply the well-known identity

$$\lim_{\substack{x_1, \dots, x_n \rightarrow a, \\ y_1, \dots, y_n \rightarrow b}} \frac{\det_{1 \leq i, j \leq n} (f(x_i, y_j))}{\prod_{1 \leq i < j \leq n} (x_i - x_j)(y_i - y_j)} = \frac{1}{\prod_{j=1}^n (j-1)!^2} \det_{1 \leq i, j \leq n} \left(\frac{\partial^{i+j-2} f(a, b)}{\partial x^{i-1} \partial y^{j-1}} \right).$$

Simplifying the prefactor using (8.8), we obtain the desired result. $\square$

Proof of Lemma 9.6. In Lemma 9.7, replace x by $x + 2\zeta + 1$ and y by $y + \zeta/(\zeta + 2)$ inside the determinant, and then replace ζ by $\zeta/2$. Note that

$$\frac{F(x + \zeta + 1, y + \zeta/(\zeta + 4), \zeta + 1; \zeta/2)}{G(x + \zeta + 1, y + \zeta/(\zeta + 4); \zeta/2)} = \frac{(\zeta + 2)(\zeta + 4)A(x, y, \zeta)}{B(x, y, \zeta)}, \quad (9.6)$$

where $A, B \in \mathbb{Z}[x, y, \zeta]$ and $B(0, 0, \zeta) = \zeta^2(\zeta + 2)^2$. It follows that the Taylor expansion of (9.6) at $x = y = 0$ has the form

$$\frac{(\zeta + 4)}{\zeta^2(\zeta + 2)} \sum_{k, l=0}^{\infty} \frac{C_{kl}(\zeta)}{(\zeta^2(\zeta + 2)^2)^{k+l}} x^k y^l,$$

where $C_{kl} \in \mathbb{Z}[\zeta]$. Using (8.2), Lemma 9.7 can then be written

$$2^{\left[\frac{n(n+2)}{2}\right]} p_n(\zeta/2) = \frac{(-1)^{\binom{n+1}{2}} \det_{1 \leq i, j \leq n} (C_{i-1, j-1}(\zeta))}{\zeta^{n^2} (\zeta + 2)^{n(n-1)} (\zeta + 1)^{\left[\frac{n^2}{4}\right]} (\zeta + 4)^{n(n-1) + \left[\frac{(n-1)^2}{4}\right]}}.$$

The right-hand side is a quotient of two polynomials with integer coefficients. Since the denominator is monic, the left-hand side has integer coefficients. $\square$

Proof of Proposition 3.4. Let n be even, and write (3.4) in the form

$$\begin{aligned} \sum_{k=0}^{n(n+1)/2} 2^{k - \frac{n(n+2)}{2}} b_k \zeta^k &= \sum_{k=0}^{[n(n+2)/12]} \frac{c_k}{2^k} \zeta^k (\zeta + 1)^{4k} (\zeta^2 + 4\zeta + 1)^{\frac{n(n+2)}{4} - 3k} \\ &- \sum_{k=0}^{[(n-2)(n+4)/12]} \frac{d_k}{2^k} \zeta^{k+1} (\zeta + 1)^{4k+3} (\zeta^2 + 4\zeta + 1)^{\frac{(n-2)(n+4)}{4} - 3k}. \end{aligned} \quad (9.7)$$

Here, b_k are coefficients of the polynomial $2^{\frac{n(n+2)}{2}} p_n(\zeta/2)$, which are integer by Lemma 9.6. The numbers c_k are coefficients of $\tilde{q}_{n+1}$, which are also integers, see §5.

We need to show that d_k are integers. To this end, let $0 \leq m \leq [(n-2)(n+4)/12]$, pick out the coefficient of $\zeta^{\frac{n(n+2)}{2}-m}$ on both sides of (9.7), and multiply the result by 2^m . This leads to a triangular system of the form

$$c_m - d_m + \sum_{j < m} (\mathbb{Z}c_j + \mathbb{Z}d_j) = \begin{cases} 0, & 0 \leq m < \frac{n}{2}, \\ b_{\frac{n(n+2)}{2}-m}, & \frac{n}{2} \leq m \leq \left[\frac{(n-2)(n+4)}{12}\right]. \end{cases}$$

By induction on m , it follows that each d_m is an integer. $\square$

Proposition 3.4 implies an integrality result for the coefficients of p_n . Although it seems to be far from sharp, it may still have some interest.

Corollary 9.8. *Let $p_n(\zeta) = \sum_{k=0}^{n(n+1)/2} a_k \zeta^k$. Then, $2^{\mu_k} a_k \in \mathbb{Z}$, where*

$$\mu_k = \begin{cases} \min \left(\left\lceil \frac{n(n+2)}{12} \right\rceil, k, \frac{n(n+2)}{2} - k \right), & n \text{ even}, \\ \min \left(\left\lceil \frac{(n+1)^2}{12} \right\rceil, k, \frac{(n+1)^2}{2} - k \right), & n \text{ odd}. \end{cases}$$

Proof. When n is even, this follows easily from (9.7), using that $2^{k-\frac{n(n+2)}{2}} b_k = a_k$ and that c_k and d_k are integers. For odd n , the proof is similar. $\square$

10. THERMODYNAMIC LIMIT

In this Section, we give a formal derivation of our conjectured expression (3.12) for the free energy. Essentially, we assume that the free energy can be expressed in terms of a formal power series in the variable ζ , and then deduce (3.12) by formally taking $n \rightarrow \infty$ in Proposition 8.3. To turn this into a rigorous proof would require strong analyticity assumptions that seem difficult to verify a priori. However, one can observe that (3.12) gives the correct result not only as $\zeta \rightarrow 0$, but also at $\zeta = 1$ (because of (2.5) and (3.14)) and as $\zeta \rightarrow \infty$ (since it has the correct behaviour under $\zeta \mapsto \zeta^{-1}$). It also seems to agree with numerical experiment.

We will assume that

$$\lim_{n \rightarrow \infty} \frac{\log(p_n(\zeta))}{n^2} = \log g(\zeta) \quad (10.1)$$

exists in the algebra of formal power series in ζ . We also assume that the convergence is regular in the sense that if we factor

$$p_n(\zeta) = g(\zeta)^{n^2} \phi_n(\zeta), \quad (10.2)$$

then

$$\lim_{n \rightarrow \infty} \frac{\phi_n(\zeta)^2}{\phi_{n+2}(\zeta)\phi_{n-2}(\zeta)} = 1, \quad (10.3)$$

still in the sense of formal power series. As a weak motivation for this assumption, we note that, by Proposition 8.9 and (3.14), $\phi_n(1) \sim C n^\alpha \beta^n$, where C is different for even and odd n . Thus, (10.3) holds pointwise at $\zeta = 1$, whereas the limit

$$\lim_{n \rightarrow \infty} \frac{\phi_n(\zeta)^2}{\phi_{n+1}(\zeta)\phi_{n-1}(\zeta)} \Big|_{\zeta=1}$$

does not exist.

Let us now apply Proposition 8.3, which we write in the form

$$(1 + \zeta)^2 p_{n+1}(\zeta) p_{n-1}(\zeta) = (1 + 2\zeta)^{1+\chi(n \text{ even})} \left(1 + \frac{\zeta}{2}\right)^{1+\chi(n \text{ odd})} p_n(\zeta)^2 + \mathcal{O}(\zeta^{n+1}),$$

where $\mathcal{O}(\zeta^{n+1})$ is a polynomial divisible by ζ^{n+1} . Iterating this gives

$$(1 + \zeta)^8 p_{n+2}(\zeta) p_{n-2}(\zeta) = (1 + 2\zeta)^6 \left(1 + \frac{\zeta}{2}\right)^6 p_n(\zeta)^2 + \mathcal{O}(\zeta^n).$$

Applying the factorization (10.2) we obtain

$$g(\zeta)^8 = \frac{(1 + 2\zeta)^6 \left(1 + \frac{\zeta}{2}\right)^6}{(1 + \zeta)^8} \cdot \frac{\phi_n(\zeta)^2}{\phi_{n+2}(\zeta) \phi_{n-2}(\zeta)} + \frac{\mathcal{O}(\zeta^n)}{(1 + \zeta)^8 g(\zeta)^{2n^2} \phi_{n+2}(\zeta) \phi_{n-2}(\zeta)}.$$

Letting $n \rightarrow \infty$ and using (10.3) we conclude that

$$g(\zeta) = \frac{(1 + 2\zeta)^{\frac{3}{4}} \left(1 + \frac{\zeta}{2}\right)^{\frac{3}{4}}}{1 + \zeta}.$$

Replacing ζ by ζ^{-1} in (10.1) then gives

$$\lim_{n \rightarrow \infty} \frac{\log(\tilde{p}_n(\zeta))}{n^2} = \log g(\zeta) \quad (10.4)$$

Finally, using (10.1) and (10.4) in Corollary 9.4 gives

$$\lim_{n \rightarrow \infty} \frac{\log Z_n^{3C}(t_0, t_1, t_2)}{n^2} = \log \left(\frac{g(\zeta)}{\eta^{1/4} \theta(\lambda^3; p^3)} \right) = \frac{1}{3} \log(t_0 t_1 t_2) + \log \left(\frac{g(\zeta)}{\eta^{1/4}} \right).$$

By Lemma 9.2, this may be written as in (3.12).

REFERENCES

- [A] G. E. Andrews, *Plane partitions. III. The weak Macdonald conjecture*, Invent. Math. 53 (1979) 193–225.
- [B1] R. J. Baxter, *Three-colorings of the square lattice: A hard squares model*, J. Math. Phys. 11 (1970) 3116–3124.
- [B2] R. J. Baxter, *Eight-vertex model in lattice statistics and one-dimensional anisotropic Heisenberg chain II. Equivalence to a generalized ice-type model*, Ann. Phys. 76 (1973) 25–47.
- [BM1] V. V. Bazhanov, V. V. Mangazeev, *Eight-vertex model and non-stationary Lamé equation*, J. Phys. A 38 (2005) L145–L153.
- [BM2] V. V. Bazhanov, V. V. Mangazeev, *The eight-vertex model and Painlevé VI*, J. Phys. A 39 (2006) 12235–12243.
- [BM3] V. V. Bazhanov, V. V. Mangazeev, *Analytic theory of the eight-vertex model*, Nuclear Phys. B 775 (2007) 225–282.
- [BM4] V. V. Bazhanov, V. V. Mangazeev, *Eight-vertex model and Painlevé VI equation. II. Eigenvector results*, 0912.2163.
- [Be] B. C. Berndt, *Ramanujan’s Notebooks, Part III*, Springer-Verlag, New York, 1991.
- [BF] P. M. Bleher, V. V. Fokin, *Exact solution of the six-vertex model with domain wall boundary conditions. Disordered phase*, Comm. Math. Phys. 268 (2006) 223–284.
- [BL1] P. Bleher, K. Liechty, *Exact solution of the six-vertex model with domain wall boundary conditions. Ferroelectric phase*, Comm. Math. Phys. 286 (2009) 777–801.
- [BL2] P. Bleher, K. Liechty, *Exact solution of the six-vertex model with domain wall boundary condition. Critical line between ferroelectric and disordered phases*, J. Stat. Phys. 134 (2009) 463–485.

- [BL3] P. Bleher, K. Liechty, *Exact solution of the six-vertex model with domain wall boundary conditions. Antiferroelectric phase*, 0904.3088.
- [D] G. Darboux, *Leçons sur la théorie générale des surfaces et les applications géométriques du calcul infinitésimal*, Deuxième partie, Gauthier-Villars, Paris, 1889.
- [FH] W. Fulton, J. Harris, *Representation Theory*, Springer-Verlag, New York, 1991.
- [H] R. Hirota, Y. Ohta, J. Satsuma, *Wronskian structure of solutions for soliton equations*, Progr. Theoret. Phys. Suppl. 94 (1988) 59–72.
- [I] A. G. Izergin, *Partition function of a six-vertex model in a finite volume*, Soviet Phys. Dokl. 32 (1987) 878–879.
- [ICK] A. G. Izergin, D. A. Coker, V. E. Korepin, *Determinant formula for the six-vertex model*, J. Phys. A 25 (1992) 4315–4334.
- [KZ] V. Korepin, P. Zinn-Justin, *Thermodynamic limit of the six-vertex model with domain wall boundary conditions*, J. Phys. A 33 (2000) 7053–7066.
- [K] G. Kuperberg, *Another proof of the alternating-sign matrix conjecture*, Internat. Math. Res. Notices 1996 (1996) 139–150.
- [LS] A. N. Leznov, M. V. Saveliev, *Theory of group representations and integration of nonlinear systems $x_{a,z\bar{z}} = \exp(kx)_a$* , Physica D 3 (1981) 62–72.
- [L] E. H. Lieb, *The residual entropy of square ice*, Phys. Rev. 162 (1967) 162–172.
- [M1] I. G. Macdonald, *Affine root systems and Dedekind's η -function*, Invent. Math. 15 (1972) 91–143.
- [M2] I. G. Macdonald, *Symmetric Functions and Hall Polynomials*, Clarendon Press, Oxford, 1979.
- [MRR] W. H. Mills, D. P. Robbins, H. Rumsey, *Alternating sign matrices and descending plane partitions*, J. Combin. Theory Ser. A 34 (1983) 340–359.
- [N] M. Noumi, *Painlevé Equations through Symmetry*, Amer. Math. Soc., Providence, 2004.
- [O] S. Okada, *Enumeration of symmetry classes of alternating sign matrices and characters of classical groups*, J. Algebraic Combin. 23 (2006) 43–69.
- [P] J. Propp, *The many faces of alternating sign matrices*, in: Discrete Models: Combinatorics, Computation, and Geometry, MIMD, Paris, 2001, pp. 043–058.
- [RS] A. V. Razumov, Yu. G. Stroganov, *Three-coloring statistical model with domain wall boundary conditions. I. Functional equations*, Theoret. and Math. Phys. 161 (2009) 1325–1339.
- [R] H. Rosengren, *An Izergin-Korepin-type identity for the 8VSOS model, with applications to alternating sign matrices*, Adv. Appl. Math. 43 (2009) 137–155.
- [RSc] H. Rosengren, M. Schlosser, *Elliptic determinant evaluations and the Macdonald identities for affine root systems*, Compositio Math. 142 (2006) 937–961.
- [S] B. Schoeneberg, *Elliptic Modular Functions*, Springer-Verlag, New York–Heidelberg, 1974.
- [St] Yu. G. Stroganov, *The Izergin-Korepin determinant at a cube root of unity*, Theoret. Math. Phys. 146 (2006) 53–62.
- [W] G. N. Watson, *Theorems stated by Ramanujan. VII: Theorems on continued fractions*, J. London Math. Soc. 4 (1929) 39–48.
- [Z1] D. Zeilberger, *Proof of the alternating sign matrix conjecture*, Electron. J. Combin. 3 (1996) Research Paper 13.
- [Z2] D. Zeilberger, *Proof of the refined alternating sign matrix conjecture*, New York J. Math. 2 (1996) 59–68.
- [Zi] P. Zinn-Justin, *Six-vertex model with domain wall boundary conditions and one-matrix model*, Phys. Rev. E 62 (2000) 3411–3418.

DEPARTMENT OF MATHEMATICAL SCIENCES, CHALMERS UNIVERSITY OF TECHNOLOGY, SE-412 96 GÖTEBORG, SWEDEN

E-mail address: hjalmar@chalmers.se

URL: <http://www.math.chalmers.se/~hjalmar>